

Relationship between human blood group antigens and haemagglutination, adhesion properties of some urinary tract infection (UTI) b  
 Hiro M. Obaid<sup>1</sup>, Hajar A. Sharif<sup>2</sup>.

## Relationship between human blood group antigens and haemagglutination, adhesion properties of some urinary tract infection (UTI) bacteria

Hiro M. Obaid<sup>1</sup>, Hajar A. Sharif<sup>2</sup>.

<sup>1</sup>Foundation of Technical Education, College of Technology/ Kirkuk, Medical Lab. Techniques dept.

<sup>2</sup>Kirkuk University, Education College For Pure Science, Biology Dept.

Received 30 March 2014 ; Accepted 22 April 2014

### Abstract

For this study a total of 135 healthy person (71 female, 64 male) were eligible. Their urine samples were assayed for adhesion test, and from 94 persons of them blood samples were taken for agglutination assay. Data from the results indicated that agglutination properties was found in *E. coli* and *Proteus* with (A<sup>+</sup>, B<sup>+</sup>, AB<sup>+</sup>, AB<sup>-</sup>, O<sup>+</sup>, O<sup>-</sup>) human erythrocytes. While *Klebsiella* and *Pseudomonas* were able to agglutinate few types of human erythrocytes. The adhesion mean of bacteria were increased significantly with age increasing. No significant differences were appeared between the types of bacteria used in the assay. Significant variation was noted between the bacterial adhesion mean to cells with six, the adhesion mean was apparently more in females than of males. The bacteria were adhered significantly more to uroepithelial cells obtained from positive blood group antigen persons, the highest rate was for AB<sup>+</sup>, A<sup>+</sup> blood groups and the lowest was for B<sup>-</sup> in both sexes for all types of bacteria used.

**Key words:** Blood groups, Haemagglutination, Adhesion, UTIs bacteria.

### العلاقة بين الفصائل الدموية للإنسان وخاصة التلازن الدموي والالتصاق لبعض أنواع بكتريا التهاب المجاري البولية

هيرو محمد عبيد<sup>١</sup>، هاجر علي شريف<sup>٢</sup>.

<sup>١</sup> هيئة التعليم التقني، الكلية التقنية/كركوك، قسم تقنية التحليلات المرضية.

<sup>٢</sup> جامعة كركوك، كلية التربية للعلوم الصرفة، قسم علوم الحياة.

#### الخلاصة

شملت هذه الدراسة ١٥٣ شخصاً اصحاء (٧١ أنثى، ٦٤ ذكر) اخذت عينات ادرار منهم لاختبار خاصية الالتصاق كما واخذت من ٩٤ شخص منهم عينات دم لاختبار التلازن الدموي البكتري. بينت نتائج الدراسة بان بكتريا *E. coli* و *Proteus mirabilis* له القابلية على تلازن كريات دم حمر الإنسان للفصائل الدموية (A<sup>+</sup>, B<sup>+</sup>, AB<sup>+</sup>, AB<sup>-</sup>, O<sup>+</sup>, O<sup>-</sup>) بينما بكتريا *Klebsiella spp*, *P. aeruginosa* كان لهما القدرة على تلازن عدد قليل من الفصائل الدموية. تأثرت قابلية الالتصاق معنويًا بالمرحلة العمرية حيث ازداد معدل الالتصاق بزيادة العمر، في حين لم يلاحظ وجود فرق معنوي بين أنواع البكتريا المختلفة المستخدمة في الدراسة. اثر الجنس معنويًا في خاصية الالتصاق البكتيري، حيث كان معدل التصاق البكتيريا للخلايا الظهارية البولية في الإناث اعلى منه في الذكور. التصاق البكتيريا كان اكبر معنويًا للخلايا التي تعود الى اشخاص ذوات فصائل دم موجبة. وكان معدل الالتصاق الاعلى لفصيلة الدم A<sup>+</sup>, AB<sup>+</sup> والمعدل الاقل كان لفصيلة B<sup>-</sup> في كلا الجنسين لكل انواع البكتيريا.

الكلمات المفتاحية: الفصائل الدموية، التلازن الدموي، الالتصاق، بكتريا المجاري البولية.

**Relationship between human blood group antigens and haemagglutination, adhesion properties of some urinary tract infection (UTI) b**

**Hiro M. Obaid<sup>1</sup>, Hajar A. Sharif<sup>2</sup>.**

**Introduction**

Bacterial urinary tract infection (UTI) is the most common infection affecting the urinary tract [1]. The majority of infections is attributed to *E.coli* which is responsible for 85% of infections, *Proteus mirabilis*, *Klebsiella*, *Pseudomonas* and *Enterococcus fecalis*, are the next most frequent [2,3]. Pathogenesis of urinary tract infection involves complex interactions between an organism, the environment and potential host [4]. Bacterial adherence to mucosa is thought to be an initial and important stage to cause urinary tract infection [5]. Among some mechanisms of bacterial adherence, the role of fimbriae and their receptors is worthy of notice, in particular, type 1 fimbria [6]. The receptors for uropathogenic bacteria are carbohydrate residues located on the surface glycolipids and glycoproteins of urothelial cells. Several of bacterial adhesions mechanisms which adhere to these urothelial cell surface antigens have been identified [7]. Pili or fimbriae are nonflagellar filamentous bacterial surface appendages composed of hydrophobic proteins. Piliated bacteria stick to surfaces, both inorganic latex particles and organic animal or plant tissues. Besides bacterial binding due to this general stickiness, specific attachment to certain hosts and tissues occurs [8-10] and is thought to be a virulence factor for bacteria colonizing or causing infection of mucous surfaces [9]. Bacterial adhesins have been classified according to the agglutination patterns resulting when bacteria bind to erythrocytes from various species. Bacteria causing agglutination of guinea pig erythrocytes inhibited in the presence of D mannose are defined as carriers of type 1 pili [11, 12]. For *E. coli* causing UTI in children, the capacity to attach to human urinary tract epithelial cells in vitro was related to the severity of the UTI produced by the strain in vivo [13].

Blood group antigens are genetically determined carbohydrate structures and may affect the availability of host cell surface receptors, the host blood group antigen status may therefore influence bacterial adherence and host susceptibility to UTIS [14,15]. Adhesion can be detected by hemagglutination reaction (essentially binding of bacteria to blood cells) and a classification system has been designed using hemagglutination. Their for the aim of the present study was to register the hemagglutination and adhesion of some UTI bacteria and its relationship with blood group antigens.

**Materials and methods**

**1-Samples collection:**

From 135 healthy person in Kirkuk city (71female, 64 male), urine samples were collected in clean cups for adhesion assay, and from 94 persons of them blood samples were collected in EDTA containing test tubes for agglutination assay. The collected blood was examined for viral and bacterial tests to ensure the clearance of any organisms at General Kirkuk Hospital.

**2- Bacterial isolates.**

*E. coli*, *Proteus mirabilis*, *Klebseilla spp.*, *Pseudomonas aeruginosa* were obtained from culturing urine of patient who were suffering from UTI in Azadi Teaching Hospital. The bacteria were isolated and identified by using routine colony characters, staining and biochemical methods [16].

**3- Bacterial suspension.**

Two to three colonies from each bacteria were suspended in normal saline and washed three times, finally the bacteria was diluted in normal saline to a concentration of  $1.5 \times 10^8$  cell/ml by calibrating it with Macferland tube number 4 [9].

**Relationship between human blood group antigens and  
haemagglutination, adhesion properties of some urinary tract infection (UTI) b  
Hiro M. Obaid<sup>1</sup>, Hajar A. Sharif<sup>2</sup>.**

**4-Blood suspension:**

The anticoagulant containing blood groups ( A<sup>+</sup>, B<sup>+</sup>, AB<sup>+</sup>, O<sup>+</sup>, A<sup>-</sup>, B<sup>-</sup>, AB<sup>-</sup>, O<sup>-</sup> ) were centrifuged to separate the blood cells. The RBC washed three times with normal saline, then suspended in saline to a concentration of 1% v/v.

**5-Uro epithelial cells collection:**

Thirty ml of urine from the females under the study were collected in clean, dry container, then centrifuged at 2000 rpm for 5 minutes and washed thrice with saline and finally suspended in 8 ml saline. The epithelial cells in this suspension were transferred on to the surface of cover slips by adding the epithelial cells suspension in a petri dish containing a filter paper No.1 and allowing the cover slips to attached to the filter paper for 10 minutes then dried [9, 10].

**6-Adherence assay of epithelial cells:**

For each experiment the air dried cover slips containing the uroepithelial or buccal cells were incubated with either 1 ml of bacterial suspension or 1 ml PBS (negative control), for 1 hour at 37°C. After incubation, the suspension was washed four times (250g, 10 min) with PBS to remove any unattached bacteria. Then fixed for 15 min in methanol, washed with saline, stained by 3% Geimsa stain for 20 minutes and washed with saline to remove excess stain. The number of bacteria adhering to each of the first 50 epithelial cells was counted with oil immersion lens light microscopy (1,000×). Epithelial cells that overlapped other cells were excluded from evaluation. The mean number of bacteria per cell was calculated. All experiments were performed in duplicate [9,10].

**7-Hemagglutination assay:-**

This test was performed by mixing the bacterial suspension with the blood suspensions on a glass slide (20µl for each suspension). Agglutination was read after 3 minutes and graded as positive or negative [11,12]. The strength of the agglutination was determined further by serial two fold dilutions in saline of the bacterial suspensions with a microtiter plate. After addition of erythrocytes to each well and incubation for 2 hours at 37c°, the agglutination titer was determined, a small pellet of erythrocytes at the bottom were considered negative, and those containing an even sheet of erythrocytes across the well considered positive [11,12].

**8- Statistical analysis:** Statistical analysis was performed using Chi-square test which was used to compare categorical variables. A p-value less than 0.05 were considered significant.

**Results**

Data from the results indicated that agglutination properties was found in *E. coli* and *Proteus* with (A<sup>+</sup>, B<sup>+</sup>, AB<sup>+</sup>, AB<sup>-</sup>, O<sup>+</sup>, O<sup>-</sup>) human erythrocytes, (table 1 ). While *Klebseilla* and *Pseudomonas* were able to agglutinate few types of human erythrocytes.



**Relationship between human blood group antigens and haemagglutination, adhesion properties of some urinary tract infection (UTI) b**

**Hiro M. Obaid<sup>1</sup>, Hajar A. Sharif<sup>2</sup>.**

**Table 1 Haemagglutination ability of the bacteria**

Human blood group	Samples No.	Bacterial type			
		<i>E. coli</i>	<i>Proteus mirabilis.</i>	<i>Klebseilla spp.</i>	<i>Pseudomonas spp.</i>
A <sup>+</sup>	18	+	+	+	+
A <sup>-</sup>	3	-	-	-	-
B <sup>+</sup>	20	+	+	-	+
B <sup>-</sup>	3	-	+	-	-
AB <sup>+</sup>	20	+	+	+	+
AB <sup>-</sup>	3	+	+	-	-
O <sup>+</sup>	20	+	+	-	-
O <sup>-</sup>	7	+	-	-	-
total	94				

+ = Positive haemagglutination to human erythrocyte, - = Negative haemagglutination to human erythrocyte

Table 2 shows the number of female and male samples according to blood group. In table 3,4 its clear that the adhesion mean of bacteria were increased significantly with age increasing. No significant differences were appeared between the types of bacteria used in the assay. Significant variation was noted between the bacterial adhesion mean to cells with six, the adhesion mean was apparently more in females than of males.

**Table 2 Number of urine samples used in adhesion assay**

Blood group	A <sup>+</sup>	A <sup>-</sup>	B <sup>+</sup>	B <sup>-</sup>	AB <sup>+</sup>	AB <sup>-</sup>	O <sup>+</sup>	O <sup>-</sup>	Total number
Number of female urine samples	7	9	8	8	10	7	13	9	71
Number of male urine samples	10	6	12	6	5	11	9	5	64
Total number	17	15	20	14	15	18	22	14	135

**Table 3 Bacterial adhesion mean to female uroepithelial cells in relation to age group**

Age group in years	Bacterial adhesion mean			
	<i>E. coli</i>	<i>Proteus mirabilis.</i>	<i>Klebseilla spp.</i>	<i>Pseudomonas spp.</i>
1-9	17.3	10.9	12.1	12.5
10->19	20.1	25.5	15.6	19.7
19-29	49.5	39.8	25.8	39.6
30-39	45.8	41.2	23.3	25.2
40-49	40.3	40.9	29.2	37.4
50-59	43.2	42.2	30.0	40.4
60-80	48.8	44.4	29.9	42.6

**Relationship between human blood group antigens and haemagglutination, adhesion properties of some urinary tract infection (UTI) b**  
**Hiro M. Obaid<sup>1</sup>, Hajar A. Sharif<sup>2</sup>.**

**Table 4 Bacterial adhesion mean to male uroepithelial cells in relation to age group**

Age group in years	Bacterial adhesion mean			
	<i>E. coli</i>	<i>Proteus mirabilis.</i>	<i>Klebseilla spp.</i>	<i>Pseudomonas spp.</i>
1-9	10.6	8.5	5.9	9.3
10->19	12.8	10.1	6.6	9.7
19-29	20.4	10.8	5.5	10.8
30-39	22.3	14.2	10.3	12.6
40-49	29.1	22.6	16.8	19.3
50-59	35.5	30.8	17.7	23.6
60-80	39.6	32.4	21.9	28.5

The bacteria were adhered significantly more to uroepithelial cells obtained from positive blood group antigen persons, the highest rate was for AB<sup>+</sup>, A<sup>+</sup> blood groups. The lowest was for B<sup>-</sup> In both sexes for all types of bacteria used, with significant differences between males and females. Being the females more than males Table 5,6.

**Table 5 Bacterial adhesion mean to female uroepithelial cells in relation to blood group**

Type of bacteria	Bacterial adhesion mean in each blood group							
	A <sup>+</sup>	A <sup>-</sup>	B <sup>+</sup>	B <sup>-</sup>	AB <sup>+</sup>	AB <sup>-</sup>	O <sup>+</sup>	O <sup>-</sup>
<i>E. coli</i>	39.6	24.4	25.2	10.3	44.2	20.9	30.8	24.8
<i>Proteus mirabilis</i>	35.3	30.9	23.6	10.9	40.8	22.6	30.1	20.7
<i>Klebseilla spp.</i>	19.9	13.5	12.8	8.8	19.3	12.6	14.2	10.7
<i>P. aeruginosa.</i>	23.5	17.2	21.1	16.4	29.4	21.9	20.8	16.5

**Table 6 Bacterial adhesion mean to male uroepithelial cells in relation to blood group**

Type of bacteria	Bacterial adhesion mean in each blood group							
	A <sup>+</sup>	A <sup>-</sup>	B <sup>+</sup>	B <sup>-</sup>	AB <sup>+</sup>	AB <sup>-</sup>	O <sup>+</sup>	O <sup>-</sup>
<i>E. coli</i>	26.1	21.3	20.5	19.9	33.6	27.6	25.2	16.8
<i>Proteus mirabilis</i>	29.2	21.1	20.8	16.8	32.3	20.9	25.7	17.7
<i>Klebseilla spp.</i>	12.6	10.2	9.7	6.6	14.5	11.6	12.8	9.9
<i>P. aeruginosa.</i>	16.2	11.1	17.3	14.6	18.9	14.4	27.7	10.8

**Relationship between human blood group antigens and haemagglutination, adhesion properties of some urinary tract infection (UTI) b**

**Hiro M. Obaid<sup>1</sup>, Hajar A. Sharif<sup>2</sup>.**

**Discussion**

The results indicated that agglutination properties were found in *E. coli* and *Proteus mirabilis* with (A<sup>+</sup>, B<sup>+</sup>, AB<sup>+</sup>, AB<sup>-</sup>, O<sup>+</sup>, O<sup>-</sup>) human erythrocytes. While *Klebsiella* and *Pseudomonas* were able to agglutinate few types of human erythrocytes. This results was consistent with those of [17, 18] were they concluded that 261 of 453 strains of *E.coli* can agglutinate human erythrocytes and 91 of 148 strains of *proteus* could agglutinate human and some animals erythrocytes. The bacterial surface antigen (s) mediating mannose resistant hemagglutination of human erythrocytes may be one factor selecting for *E.coli* agglutinating ability [17]. *P. mirabilis* may express a variety of adherence organells which have been characterized by their ability to agglutinate various erythrocyte types [19].

Bacterial adhesion means was greater in females than those of males in all bacteria used this may due to that the women has UTIS more than the men [1] and this may leads to alteration and increased bacterial adherence, this has been described in a studies on vaginal and buccal epithelial cells isolated from patients (women) with recurrent UTIS [20]. In addition of some special factors acting in females may influence bacterial adhesions like estrogen hormans levels. An increased bacterial adherence was occurred during the menstrual cycle and early pregnancy [21].

Adhesion mean of the bacteria to the uroepithilial in the present study was differed according to the blood group antigens. Blood group antigens are carbohydrate structures bound to membranes lipids or proteins are integral structure in uroepithelial cell membranes. This affect the ability of host cell surface receptors, the host blood group antigen status may therefore influence bacterial adherence and host susceptbility to UTIS [15]. It has been suggested that bacterial adherence may also be affected by the blood group antigens, which are found on the surface of uroepithelial cells. Individuals with the Lewis blood group phenotype Le(a<sub>-</sub>b<sub>-</sub>) secrete Leb and A, B, or H substances in their saliva and plasma and are called "secretors," whereas nonsecretors with the Le(a<sub>-</sub>b<sub>-</sub>) phenotype do have Lea antigens in their secretions but not A, B, or H substances. Several studies have shown a correlation between the Lewis blood group phenotypes and recurrent UTIs in adult women [14, 15]. ABH non-secretors are at a greater risk for recurrent urinary tract infections (UTI) and are much more likely to develop renal scars. This susceptbility is even greater among the Lewis negative subset (Le (ab<sub>-</sub>)).The ABH secretor phenotype conveys a measure of protection; cutting the risk of recurrent UTI by greater than 50% and dramatically decreasing the likelihood that renal scars will develop. ABH non-secretors appear to be at extra risk for recurrent urinary tract infections. In one study of women with recurrent UTI, 29 % of the women were the Lewis (a<sup>+</sup> b<sup>-</sup>) non-secretor phenotype, while another 26% of the women were Lewis (a<sup>-</sup> b<sup>-</sup>) recessive phenotype. When the women with ABH non-secretor and recessive phenotypes were combined and considered collectively, the odds ratio (an estimate of relative risk of urinary tract infection) for those without the secretor phenotype (Lewis (a<sup>-</sup> b<sup>+</sup>)) was 3. A form of synergy also appears to exist between UTI risk, secretor status and the lack of ability to create anti-B isohemagglutinin. Essentially, blood group B and AB and the non-secretor phenotype seem to work together to increase the relative risk of UTI among these women [22].



**Relationship between human blood group antigens and  
haemagglutination, adhesion properties of some urinary tract infection (UTI) b**

**Hiro M. Obaid<sup>1</sup>, Hajar A. Sharif<sup>2</sup>.**

**References**

1. Fioriti D, Penta M, Mischitelli, *et al.* : Interstitial cystitis and infections agents. *J. Pharma.* 2005, 18 (4) : 799–804.
2. Dromigny JA, Nabeth P, Juergens BA, Perrier-Gros-Claude JD. Risk factors for antibiotic-resistance *Escherichia coli* urinary tract infection in Dakar; Senegal. *J. Antimicrob. Chemother.* 2005, 56: 236-239.
3. Mahmood MA. Prevalence and Antimicrobial Susceptibility of Pathogens in Urinary Tract Infections. *Journal of Al-Nahrain University* .2011,14 (4):46-152.
4. Anderson GG, Martin SM and Hultgren SJ. Host subversion by formation of intracellular bacterial communities in the urinary tract. *Microbes Infect* ,2004,6(12):1094-10.
5. Mulvey MA. Adhesion and entry of uropathogenic *Escherichia coli*. *Cell Microbiol*, 2002,4(5), 257-71.
6. Richard AH, William HD, Michael DT, *et al.* Rote of Type 1 Fimbria- and P Fimbria-specific Adherence in colonization of the Neurogenic Human Bladder by *E. coli*. *American Soci. Micro.* 2002: 70 (11): 6481–6484
7. Lau YE, Rozek A, Scott MG, Goosney DL, Davidson DJ and Hancock R E. Interaction and cellular localization of the human host defense peptide LL-37 with lung epithelial cells', *Infect Immun*,2005, 73(1): 583-91.
8. Wu XR, Sun TT, Medina JJ: In vitro binding of type 1-fimbriated *Escherichia coli* to uroplakins Ia and Ib: relation to urinary tract infections. *Proc Natl Acad Sci U S A* ,1996, 93:9630–9635.
9. Hull RA, Donovan WH, Terzo MD. Role of type 1- fimbria and P-fimbria specific adherence in colonization of neurogenic human bladder by *Escherichia coli* .IAI. infection and immunity , 2002, 70 (11):6481-6484.
10. Ghenghesh K S, Elkateb E, Berbash N, Nada R A, Salwa F, Rahouma AA, Seif- nasser N, Elkhatabroun MA, Belresh T and John D : Diabetic patients in Libya: Virulence factors and phylogenetic group of *Escherichia coli* isolates. *J. Med. Microbiol.*, 2009, 58: 1006-1014.
11. Andreu A, AE Stapletou, CL Fennell, *et al.* Hemegglutination, adherence and surface properties of vaginal lactobacillus sp. *J. Infee. Dis.* 1995, 171 : 1237–1243.
12. Graziela B, Marilda C. Evaluation of Adherence, Hemagglutination, and presences of Gens codifying for virulence factors of *Acidobacter baumannii* causing urinary tract infection. *Riode Janerio.* 2004 : 99 (8) : 839–844.
13. Svanborg Eden C, Jodal U: Attachment of *Escherichia coli* to urinary sediment epithelial cells from urinary tract infection prone and healthy children. *Infect Immun.* 1997,26:837–840.
14. Kinane DF, Blackwell CC, Brettle RP, Weir DM, Winstanley FP, Elton RA: ABO blood group, secretor state, and susceptibility to recurrent urinary tract infection in women. *Br Med J.* 1982, 285:7–9.
15. Sheinfeld J, Schaeffer AJ, Cordon Cardo C, Rogatko A, Fair WR: Association of the Lewis blood-group phenotype with recurrent urinary tract infections in women. *N Engl J Med.* 1989, 320:773–777.
16. Colle JG, Miles RS, Watt B. Tests for the identification of bacteria. In:Mackie and MacCartney Practical Microbiology, 14th edn. New York: Churchill Livingstone Inc., 1996, 131- 49.

**Relationship between human blood group antigens and haemagglutination, adhesion properties of some urinary tract infection (UTI) b**

**Hiro M. Obaid<sup>1</sup>, Hajar A. Sharif<sup>2</sup>.**

17. Hagberg IARS, Timok U *et al.* Adhesion, Hemagglutination and virulence of *Escherichia coli* causing urinary tract infections. *Infec. Immun.* 1991, 31(1): 561–570.
18. Mis hra M, Thakar YS, Pathak AA, Haemagglutination, haemolysin production and serum resistance of proteus and related species isolated from clinical sources. *Official puplicatiuon of indion association of medical microbio.* 2001: I2: 5–11.
19. Xin Li, David E. Johnson and Harry L.T. Molbley Requirement of Mr PH for mannose. Resistant proteus. Like Fimbria. Mediated Hemagglutination by proteus mirabilis. *Infect. Immun.* 1999. 67(6): 2822–2833.
20. Schaeffer AJ, Jones JM and Dunn JK. Association of vitro *Escherichia coli* adherence to vaginal and buccal epithelial cells with susceptibility of women to recurrent urinary-tract infections', *N Engl J Med*, 1981, 304(18), 1062-6.
21. Haspels AA, Luisi M and Kicovic PM. Endocrinological and clinical investigations in post-menopausal women following administration of vaginal cream containing oestriol. *Maturitas*, 1981, 3(3-4), 321-7.
22. NAP 2007 Technical Reference Manual, ABH Secretor Test. (Cost effective salivary test to determine ABH secretor status) [TE002]

