

## On The Product Of Positively Defined Operators

Mahmood Kamil Shihab

## On The Product of Positively Defined Operators

Mahmood Kamil Shihab

Department Of Mathematics- College Of Science - Diyala University – Iraq

Received: 27 May 2015

Accepted: 26 April 2016

**Abstract**

The main result is to prove that the product of two positively defined operators is positively defined if and only if it is normal. In general, the normality is required cannot be dropped.

**Keywords:** Positively defined operators, Normal Operators, self-adjoint operators, Hilbert space.

**حول حاصل ضرب المؤثرات المعرفة تعريفا ايجابيا**

محمود كامل شهاب

قسم الرياضيات - كلية العلوم جامعة ديالى - العراق

**الخلاصة**

النتيجة الرئيسية هي برهنة أن حاصل ضرب مؤثرين معرفين تعريفا ايجابيا يكون مؤثرا معرفا تعريفا ايجابيا إذا فقط إذا كان ناتج حاصل ضربهما مؤثرا طبيعيا. على العموم ناتج حاصل ضرب المؤثرين مؤثر طبيعي أمر ضروري لا يمكن إسقاطه. **الكلمات المفتاحية:** المؤثرات المعرفة تعريفا ايجابيا، المؤثرات الطبيعية، مؤثر ذاتية المرافق، فضاء هلبرت.

**Introduction**

In 2002 Hichem M. Mortad communicated with Joseph A. Ball [2] proved the fact that if we have two self-adjoint operators (bounded or not) and if their product is normal, then it is self-adjoint provided a certain condition  $\sigma(A) \cap \sigma(-A) \subseteq \{0\}$  satisfied, where  $A$  is one of two self-

## On The Product Of Positively Defined Operators

Mahmood Kamil Shihab

adjoint operators. However, nothing was said about the product of two positively defined operators, is positively defined or not? In this paper we answer this question positively, and we show that the normality is necessary for this case, we give a counter example.

We also note that one can prove this result without the condition  $\sigma(A) \cap \sigma(-A) \subseteq \{0\}$  we use the positivity of operators instead of this condition. We also use the famous Fuglede-Putnam Theorem.

### Results

We recall Hichem M. Mortad Theorem [2]

**Theorem 1.** *Let  $H$  and  $K$  be two bounded self-adjoint operators. Let  $K$  satisfies the condition  $\sigma(A) \cap \sigma(-A) \subseteq \{0\}$  If  $HK$  is normal then it is self-adjoint.*

We also recall the Fuglede-Putnam Theorem [3]

**Theorem 2.** *For normal operators  $N_1$  and  $N_2$  and arbitrary operator  $A$ , if  $AN_1 = N_2 A$  then  $AN_1^* = N_2^* A$ .*

**Definition 1.** *A bounded operator  $A$  is said to be positively defined if  $A$  is self-adjoint and  $\langle Ax, x \rangle \geq 0$  for all  $x \in H$ , where  $H$  is a Hilbert space.*

We also recall the following Theorems [1]:

The following theorem with proof can be found in [1, Theorem 4.6.14]

**Theorem 3.** *Every positively defined operator  $A$  has a unique positively defined square root  $B$ . Moreover,  $B$  commutes with every operator commuting with  $A$ .*

The following theorem with proof can be found in [1, Theorem 4.6.9]

**Theorem 4.** *Product of two commuting positively defined operators is positively defined.*

We present here the proof of known proposition.

**Proposition 1.** *Let  $A$  and  $B$  be two self-adjoint operators, then  $AB$  is normal if and only if  $BA$  is normal.*

#### **Proof.**

Let  $AB$  be normal that is

$$AB(AB)^* = (AB)^* AB$$

$$\begin{aligned} BA(BA)^* &= BAA^* B^* \\ &= (AB)^* AB \end{aligned}$$

On The Product Of Positively Defined Operators

Mahmood Kamil Shihab

$$\begin{aligned}
 &= AB(AB)^* \\
 &= ABB^* A^* \\
 &= (BA)^* BA.
 \end{aligned}$$

Conversely, similarly we get

$$AB(AB)^* = (AB)^* AB.$$

**Proposition 2.** Let  $H$  be a Hilbert space, if  $A$  is a positively defined in  $H$  then  $A^2$  is also a positively defined in  $H$ .

The main result:

**Theorem 5.** Let  $H$  be a Hilbert space, let  $A$  and  $B$  be two positively defined operators on  $H$ ,  $AB$  is positively defined operator if and only if  $AB$  is normal.

**Proof.**

The proof of  $AB$  is normal when  $AB$  is positively defined is obvious.

Conversely, let  $AB$  be normal, by **Proposition 1** also  $BA$  is normal, corresponding to **Theorem 2** let

$$N_1 = BA \quad \text{and} \quad N_2 = AB$$

and  $A$  is an arbitrary bounded operator so we have

$$ABA = ABA$$

then

$$A(BA)^* = (AB)^* A$$

$$AA^* B^* = B^* A^* A$$

$$AAB = BAA$$

$$A^2 B = BA^2.$$

Since  $A^2$  commutes with  $B$  and  $A$  is positively defined **Proposition 2** implies  $A^2$  is also positively defined, and so by **Theorem 3** we have

$$AB = BA.$$

So by **Theorem 4** we get  $AB$  is positively defined.

3- Counter Example

## On The Product Of Positively Defined Operators

Mahmood Kamil Shihab

The normality of  $AB$  in **Theorem 5** is necessary for positivity of  $AB$ . If  $AB$  is not normal then  $AB$  is not always positively defined, we give the following counter example:

**Example 1.** Let

$$A = \begin{bmatrix} 1 & i \\ -i & 1 \end{bmatrix} \quad B = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$$

it easy to check that both  $A$  and  $B$  are positively defined, and  $AB$  is not normal. Also it easy to check that  $AB$  is not self-adjoint and so it is not positively defined.

### References

1. Loenath Debnath and Piotr Mikusinski: *Introduction to Hilbert Spaces with Applications*, Elsevier Academic Press, 2005 (3rd edition).
2. Hichem M. Mortad (communicated by Joseph A. Ball): An Application of The Putnam-Fuglede Theorem to normal products of self-adjoint operators, American Mathematical Society, 131/10 (2003) 3135-3141.
3. C.R. Putnam: On The Operator In Hilbert Space, Amer. J. Math., 73 (1951) 357-362.
4. B. Fugled: A Commutativity Theorem For Normal Operators, National Acad. Sci., Vol 36 (1950) 35-40.
5. W. Rudin: *Functional Analysis*, McGraw Hill Newyourk, 1991 (2nd edition).