

**Phytochemical Analysis and Inhibitory Effect of *Citrus aurantium* L.****(Bitter Orange) Leaves on some Bacterial Isolates *in vitro***

Khudhair A. M. Abed AL Ani

Tareq R. Minnat

Osama K. Jalyl

**Phytochemical Analysis and Inhibitory Effect of *Citrus aurantium* L.****(Bitter Orange) Leaves on some Bacterial Isolates *in vitro***Khudhair A. M. Abed AL Ani<sup>1</sup>, Tareq R. Minnat<sup>2</sup> and Osama K. Jalyl<sup>3</sup><sup>1</sup> Physiology and Pharmacology Dept., College of Veterinary Medicine, Diyala University.<sup>2</sup> Medicine Dept., College of Veterinary Medicine, Diyala University.<sup>3</sup> Microbiology Dept., College of Veterinary Medicine, Diyala University.**Received: 15 August 2016****Accepted: 27 September 2016****Abstract**

This study has been done in order to investigate the phytochemicals constitution and determine *in vitro* antimicrobial activity of aqueous and methanolic *Citrus aurantium* L. leaves extract against bacteria. The organic solvent extraction done by used Soxhlet apparatus, solvents used were methanol (80%), while aqueous extract was prepared by using magnetic stirrer. The phytochemical analysis of plant extract was performed using precipitation and coloration methods. Antibacterial activity of methanolic and aqueous of *Citrus aurantium* L. leaves extract determined by disc diffusion and agar well diffusion method. The different concentrations of aqueous and methanolic *Citrus aurantium* L. leaves extract exhibited potent antibacterial activity against selected bacteria. The phytochemical screening of aqueous and methanolic *Citrus aurantium* L. leaves extract revealed the presence of steroids, flavonoids, saponins, alkaloids, phenols and tannins, while the glycosides detectable in the methanolic extract only, and the terpenoids not detectable. Both aqueous and methanolic *Citrus aurantium* L. leaves extract at concentrations (5% and 10%) showed considerable antibacterial activity against (*Staphylococcus epidermidis* and *Staphylococcus aureus*, *E. coli*, *Pseudomonas aeruginosa* and *Proteus mirabilis*). To conclude, the antimicrobial activity of the aqueous and methanolic extracts of *Citrus aurantium* L. leaves was efficiency against some bacteria strains.

**Keywords:** - *Citrus aurantium* L., Phytochemical, Antibacterial, Phenols, Alkaloids.

Phytochemical Analysis and Inhibitory Effect of *Citrus aurantium* L.(Bitter Orange) Leaves on some Bacterial Isolates *in vitro*

Khudhair A. M. Abed AL Ani

Tareq R. Minnat

Osama K. Jalyl

## تحليل مركبات الايض الثانوية والتأثير التثبيطي لأوراق نبات النارنج على بعض انواع البكتيريا خارج الجسم الحي

خضير عبد الرحمن محمود ، طارق رفعت منت<sup>2</sup> و اسامة غازي جليل<sup>3</sup><sup>1</sup> فرع الفلسفة والأدوية – كلية الطب البيطري – جامعة ديالى<sup>2</sup> فرع الطب الباطني – كلية الطب البيطري – جامعة ديالى<sup>3</sup> فرع الأحياء المجهرية – كلية الطب البيطري – جامعة ديالى

### الخلاصة

أجريت هذه الدراسة لغرض الكشف عن وجود مركبات الايض الثانوية وتقييم الفعالية ضد البكتيريا في المختبر لكل من المستخلص المائي والكحولي لأوراق النارنج. استخدم جهاز السوكسلت لأستخلاص بالمذيبات العضوية فقد أستخدم الميثانول بنسبة (80%). بينما حضر المستخلص المائي بواسطة استخدام جهاز الرجاج المغناطيسي. تم الكشف عن مركبات الايض الثانوية باستخدام الطرائق اللونية وطريقة الترسيب . اما الفعالية المضادة للبكتيريا لكل من المستخلص المائي والكحولي لأوراق النارنج فقد قيمت بطريقة الأنتشار في الأكار بواسطة الحفر وطريقة الاقراص المشربة. حيث أظهرت التراكيز المختلفة لكل من المستخلص المائي والكحولي لأوراق النارنج فعالية ضد البكتيريا. التحليل النوعي للمكونات الايض الثانوية لكل من المستخلصين المائي والكحولي لأوراق النارنج أظهرت وجود كل من الستيرويدات والفلافونيدات والصابونينات والقلويدات والفينولات والعصيات بينما الكلايكوسيدات حددت في المستخلص الكحولي فقط و التيربينويدات لم تحدد في المستخلص المائي والكحولي . كلا المستخلصان المائي والكحولي بتركيز (5% ، 10%) أظهرت فعالية كبيرة ضد (*Staphylococcus epidermidis* and *Staphylococcus aureus*, *E. coli*, *Pseudomonas aeruginosa* and *Proteus mirabilis*). الاستنتاج ، الفعالية المضادة للبكتيريا لكل من المستخلص المائي والكحولي لأوراق النارنج كانت كفوءة ضد بعض العتري البكتيرية.

**الكلمات المفتاحية:** - النارنج , مركبات الايض الثانوية , مضاد للبكتيريا , الفينولات , القلويدات.

### Introduction

*Citrus aurantium* L., previously known as *Fructus aurantii*, is the botanical name of a plant known as green orange, bitter orange or sour orange.<sup>[1,2]</sup> The extract of the immature fruit or peel of *C. aurantium* L. has been widely used in weight loss dietary supplements and in sports performance products<sup>[3]</sup>. In addition, *Citrus aurantium* L. is among the species that have been used for medical purposes on account of the numerous bioactive compounds that it contains such as phenolic, essential oils, vitamins and flavonoids. *Citrus aurantium* L. used for

Phytochemical Analysis and Inhibitory Effect of *Citrus aurantium* L.  
(Bitter Orange) Leaves on some Bacterial Isolates *in vitro*

Khudhair A. M. Abed AL Ani

Tareq R. Minnat

Osama K. Jalyl

medicinal purposes such as the leaves, peel and flowers of the plant.<sup>[4]</sup> Many active phytoconstituents such as flavonoids have a large spectrum of biological activity including antibacterial, antiviral, anticancer, antifungal and antidiabetic activities.<sup>[5, 6]</sup> The phytochemicals are also known as secondary metabolites that are derived from primary metabolites and are used as drugs.<sup>[7]</sup> The phytochemicals also contribute to the flavor, colour and other characteristics of plant parts.<sup>[8]</sup> Phytochemical analysis is benefit in determination of some active biological constituents of some vegetables and medicinal plants.<sup>[9]</sup> Phytochemicals are not essentially required for the sustenance of life but confer extra health benefits against bacteria.<sup>[10]</sup> In the last three decades, a numerous of antibiotics and other synthetic drugs were produced in the world with an aim of eradicating the microorganisms which were responsible for many diseases.<sup>[11]</sup> Drugs or antibiotics induced mutations in the genetic composition of these microorganisms rendering them resistant to several drugs or antibiotics.<sup>[12]</sup> Moreover, the side effects associated with the long-term use of the synthetic drugs may cause serious damages to many of human organs. Therefore, to overcome this limitation of synthetic drugs, researchers have shifted their focus towards medicinal plants which are recognized as rich sources of antibacterial agents and are widely used by different countries for medicinal purposes. Traditionally used medicinal plants are known to produce a variety of compounds with therapeutic properties, such as antidiabetic, antioxidant, antibacterial, anti-inflammatory, antipyretic, gastroprotective effects, etc.<sup>[13]</sup> The aim of this study was to investigate the phytochemicals constitution and determine *in vitro* antimicrobial activity of *Citrus aurantium* L. leaves extract against *Staphylococcus aureus*, *Staphylococcus epidermidis*, *E.coli*, *Pseudomonas aeruginosa* and *Proteus mirabilis*.

## Materials and Methods

### Collection of Plant

Fresh leaves of *Citrus aurantium* L. plant were collected from Baqubah, Diyala Province, Iraq. Which was identified, authenticated by Assist. Prof. (dr.) Khazal Dhbea Wadi, College of Sciences, Diyala University. The leaves were cleaned, washed then dried in room temperature at shade. The leaves were then crushed by using mechanical grinding machine.



**Phytochemical Analysis and Inhibitory Effect of *Citrus aurantium* L.  
(Bitter Orange) Leaves on some Bacterial Isolates *in vitro***

Khudhair A. M. Abed AL Ani

Tareq R. Minnat

Osama K. Jalyl

### **Preparation of Extracts**

**Aqueous Extract:** Thirty grams of leaves powder was suspended in 300 ml of distilled water. Aqueous extraction was done at 70°C for 30 minutes, after that filtering of the extracts by using Whatman filter paper No.1. Then evaporated the extracts at 45°C to form a paste and after that transferred into sterile bottles and refrigerated 4°C until use.

**Methanolic Extract:** Organic solvent extraction of the *Citrus aurantium* L. leaves powder was carried out by using methanol (80%). This was done by using Soxhlet apparatus. The extraction was carried out for 24 hours by heating temperature that kept the solvent at 50-60°C until a colorless and clear solvent appeared in the extracting chamber. After that, the extract was dried by electric oven at temperature 40-45°C until dry extract was obtained. The final extract was kept frozen at (-18°C) until use.

### **Phytochemical Analysis (Qualitative Analysis):**

Phytochemical analysis was carried out to evaluate the qualitative of chemical composition of crude extracts to identify the secondary metabolites groups such as glycosides, phenolic, compounds alkaloids, tannins, saponins, terpenoids, steroids and flavonoids by using coloration and precipitation methods. This analysis revealed the presence or absence of the phytochemicals constitution in the aqueous and methanolic extract.<sup>[14-18]</sup>

### **Microbial Isolates**

Different five clinical microbial isolates (Gram positive and Gram negative) were collected and identified by using conventional biochemical tests<sup>[19]</sup> and cultivated in pure culture, at Microbiological Laboratory/ College of Veterinary Medicine / Diyala University. These include (*Staphylococcus epidermidis*, *Staphylococcus aureus*, *E. coli*, *Pseudomonas aeruginosa* and *Proteus mirabilis*).

### **Antibacterial Activities**

#### **Agar Well Diffusion Method**

Determination of extract activities by spreading of 0.1 ml of bacterial suspension prepared according<sup>[20]</sup> which contain  $1.5 \times 10^8$  cell/ml over the surface of Muller – Hinton agar plate to obtain uniform growth, left the plate to dry for five minutes. Then the wells were prepared by using Cork borer five mm diameter. These wells filled by 50 microliter concentrated extract

**Phytochemical Analysis and Inhibitory Effect of *Citrus aurantium* L.****(Bitter Orange) Leaves on some Bacterial Isolates *in vitro*****Khudhair A. M. Abed AL Ani****Tareq R. Minnat****Osama K. Jalyl**

of either methanolic or aqueous extract according to concentration used 2.5%, 5% and 10%, the medium was left for 1 hour in laboratory condition, then incubated for 24 h at 37 °C and the inhibition zone of antibacterial activity appeared around the well were measured the diameter by centimeter scale in mm (millimeter), each treatment consists of 4 repeat. [21-23]

**Disc Diffusion Method**

Antibacterial activity of aqueous and methanolic extracts were determined by disc diffusion method on Muller –Hinton agar by (Steel and Torrie ). [24] Sterile Whatman filter disc (5 mm diameter) were made, these disc impregnated in the 50 microliter of aqueous or methanolic extract. Placed in Petri dishes according to concentrations (2.5%, 5% and 10%) for 24 hours. Inoculums containing  $1.5 \times 10^8$  CFU / ml of bacteria were spread, with sterile swab moistened with the bacterial suspension. The disc also impregnated in 50  $\mu$ L of solvent distilled water served as a standard control. [23] Standard antibiotic disc; Tetracycline (T); Streptomycin (S); Penicillin (P); Amoxicillin (AUG) for antibacterial activity tests were carried out against bacterial strains.

**Statistical Analysis**

The values are expressed as the mean  $\pm$  the standard error of the mean (M $\pm$ SEM). The values were analyzed by using one way analysis of variance ANOVA, and then the test of the least significant difference LSD applied to find the significant differences between the means of inhibitory zones . [25] The significant level of test was  $P < 0.05$ .

**Results and Discussion****Phytochemical Analysis**

The results of the preliminary phytochemical screening of aqueous and (80%) methanolic *Citrus aurantium* L. leaves extract revealed the presence of a wide range of phytochemicals constitution including steroids, tannins, alkaloids, flavonoids, saponins, phenol while the glycosides are detected only in the *Citrus aurantium* L. methanolic extract but the terpenoid don't detectable in the aqueous and methanolic extract. Table (1).

Phytochemical Analysis and Inhibitory Effect of *Citrus aurantium* L.(Bitter Orange) Leaves on some Bacterial Isolates *in vitro*

Khudhair A. M. Abed AL Ani

Tareq R. Minnat

Osama K. Jalyl

**Table 1: Phytochemical screening of aqueous and (80%) methanolic *Citrus aurantium* L. leaves extract**

No.	Test	<i>Citrus aurantium</i> L. Aqueous Extract	<i>Citrus aurantium</i> L. 80% methanolic Extract
1	Alkaloid	+	+
2	Flavonoid	+	+
3	Steroid	+	+
4	Terpenoid	—	—
5	Phenol	+	+
6	Saponins	+	+
7	Tannins	+	+
8	Glycosides	—	+

[+ = Present, — = Absent ]

In the present study, we evaluated the phytochemical compound of aqueous and methanolic extract of *Citrus aurantium* L. leaves. The analysis of phytochemicals constitution conducted on the plant extracts revealed the presence of phytochemicals constitution which are known to exhibit medicinal as well as physiological activities.<sup>[26]</sup> The citrus peels are rich in nutrients and contain many phytochemicals with strong potential for use in drug production or as food supplements.<sup>[27,28]</sup> Our results are in agreement with these assertions as a range of phytochemicals; alkaloids, tannins, flavonoids, saponins, cardiac glycosides, steroids were detected in the orange peels and seeds extracts. (Swapnil Mehta *et al.*, 2011)<sup>[29]</sup>, has reported that phytochemical analysis of *Citrus maxima* showed the presence of important classes of phytoconstituents like alkaloids, saponins and carbohydrates, this is similar to the results obtained in the present study.

**Antibacterial Activity**

The present study shown that the inhibitory zones were more at (10%) in comparison with (2.5%, 5%) of aqueous and methanolic *Citrus aurantium* L. leaves extract in each disc diffusion and well diffusion method against all bacteria species. On the other hand, the present study show significant difference at  $P < 0.05$  between susceptibility of bacteria species to 5% and 10% concentration of methanolic extract of *Citrus aurantium* L. leaves extract in well diffusion as in (Table 2). The methanolic extract of *Citrus aurantium* L. leaves extract in

Phytochemical Analysis and Inhibitory Effect of *Citrus aurantium* L.(Bitter Orange) Leaves on some Bacterial Isolates *in vitro*

Khudhair A. M. Abed AL Ani

Tareq R. Minnat

Osama K. Jalyl

well diffusion method shown inhibition zone 5% and 10% concentration more than in disc method.

**Table (2): Comparison the effect of methanolic extract of *Citrus aurantium* L. leaves extract in Well diffusion and disc diffusion methods against some bacteria species.**

Bacteria species	Methanolic extract of <i>Citrus aurantium</i> L.					
	Well diffusion			disc diffusion		
	Concentration of extract			Concentration of extract		
	2.5% M±SE (mm)	5% M±SE (mm)	10% M±SE (mm)	2.5% M±SE (mm)	5% M±SE (mm)	10% M±SE (mm)
<i>Staphylococcus aureus</i>	0.0±0.0	8 ±0.0 -b-	12.6±0.35 -c-	0±0.0	7±0.0 -b-	10.3±0.35 -c-
<i>Staphylococcus epidermidis</i>	7±0.0 -a-	10.6±0.35 -b-	12.3±0.35 -c-	0±0.0	0±0.0 -b-	10±0.33 -c-
<i>E. coli</i>	0±0.0	9±0.0 -b-	11±0.0 -c-	0±0.0	7±0.0 -b-	10±0.33 -c-
<i>Pseudomonas aeruginosa</i>	0±0.0	7±0.0 -b-	9±0.0 -c-	0±0.0	0±0.0 -b-	7±0.33 -c-
<i>Proteus mirabilis</i>	0±0.0	7.6±0.33 -b-	9.6±0.33 -c-	0±0.0	0±0.0 -b-	8±0.0 -c-

Values: Mean ± Standard error of the mean (M± S.E.M). a, b, c; significantly different level of P < 0.05  
 -a- mean the significance was between concentration 2.5% and 5%.  
 -b- mean the significance was between concentration 5% and 10%.  
 -c- mean the significance was between concentration 2.5% and 10%.

The aqueous extract of *Citrus aurantium* L. leaves in agar well diffusion method exhibit a significantly higher zones of inhibitions against gram positive bacteria (*Staphylococcus epidermidis* and *Staphylococcus aureus*) and gram negative bacteria (*E. coli*, *Pseudomonas aeruginosa* and *Proteus mirabilis*) in used at 10%. Comparison with 2.5%, 5% and 10% concentration (Table 3). On the other hand the aqueous extract of *Citrus aurantium* L. in well diffusion method shown inhibition zone 10% and 5% concentration more than in disc method.



Phytochemical Analysis and Inhibitory Effect of *Citrus aurantium* L.(Bitter Orange) Leaves on some Bacterial Isolates *in vitro*

Khudhair A. M. Abed AL Ani

Tareq R. Minnat

Osama K. Jalyl

**Table (3): Comparison the effect of aqueous extract of bitter orange in Well diffusion and disc diffusion methods against some bacteria species.**

Bacteria species	Aqueous extract of <i>Citrus aurantium</i> L.					
	Well diffusion			disc diffusion		
	Concentration of extract			Concentration of extract		
	2.5% M±SE (mm)	5% M±SE (mm)	10% M±SE (mm)	2.5% M±SE (mm)	5% M±SE (mm)	10% M±SE (mm)
<i>Staphylococcus aureus</i>	0.0±0.0	7±0.0 -b-	11.6±0.35 -c-	0.0±0.0	0±0.0 -b-	10±0.33 -c-
<i>Staphylococcus epidermidis</i>	0.0±0.0 -a-	8.6±0.35 -b-	13.6±0.35 -c-	0.0±0.0 -a-	9.6±0.33 -b-	12.3±0.35 -c-
<i>E. coli</i>	0.0±0.0	8.6±0.35 -b-	12.6±0.5 -c-	0.0±0.0	0±0.0	11±0.0
<i>Pseudomonas aeruginosa</i>	0.0±0.0	7.6±0.5 -b-	10±0.33 -c-	0.0±0.0	0±0.0	7±0.0
<i>Proteus mirabilis</i>	0.0±0.0	7±0.0 -b-	9±0.0 -c-	0.0±0.0	0±0.0	8±0.0

**Values: Mean ± Standard error of the mean (M± S.E.M). a, b, c ; significantly different level of P < 0.05**  
 -a- mean the significance was between concentration 2.5% and 5%.  
 -b- mean the significance was between concentration 5% and 10%.  
 -c- mean the significance was between concentration 2.5% and 10%.

Antibacterial efficacy of aqueous and methanolic extract of *Citrus aurantium* L. (Bitter Orange) leaves against five bacteria by agar well diffusion and disc diffusion method showed excellent inhibitory action against all tested bacteria at concentration (5%, 10%), but don't shown inhibition at concentration (2.5%) [Table 1,2]. Citrus fruit products such as *Citrus aurantium* L. act as potent antimicrobial agents against micro-organism (bacteria and fungus) because citrus fruits are rich source of flavonones and many polymethoxylated flavones, in other plants this phytoconstituents are very rare.<sup>[30]</sup>

In the study, Kirbaşlar *et al.*<sup>[31]</sup> evaluated antibacterial activity of peel oil from Citrus fruits. It was observed that peel oil of *C. sinensis* and *C. aurantium* exhibited more or less similar inhibition of Gram-positive and Gram-negative bacteria.

Bouabdelli *et al.*<sup>[32]</sup> showed that *S. aureus*, *Proteus mirabilis* and *E. coli* were sensitive to aqueous extract of *C. aurantium* L. leaves while *P. aeruginosa* was resistant to it. In addition, Gopal illustrated that extracts of *C. aurantium* L. leaves were effective against both Gram-



**Phytochemical Analysis and Inhibitory Effect of *Citrus aurantium* L.****(Bitter Orange) Leaves on some Bacterial Isolates *in vitro*****Khudhair A. M. Abed AL Ani****Tareq R. Minnat****Osama K. Jalyl**

negative and Gram-positive organisms with inhibition zone (12-14 mm). Preliminary phytochemical analysis of the *C. aurantium* L. leaf extracts showed the presence of tannins, terpenoids, sterols, alkaloids, flavonoids and carbohydrates.<sup>[33]</sup> Also Kabra, *et al.*,<sup>[34]</sup> reported that ethanolic extract of *Citrus medica* showed antibacterial activity when tested against *Klebsiella pneumoniae*, *Escherichia coli* and *Staphylococcus aureus*. Cano, *et al.*,<sup>[35]</sup> reported the higher antimicrobial activity of *Citrus paradise* due to the presence of essential oils, vitamin C and flavonoids. Dhanavade, *et al.*,<sup>[36]</sup> reported the presence of alkaloids, lactons, polyacetylene, essential oils, corydaline, pseudohypericin, protopine, hypericin and acyclic sesquiterpenes compounds from lemon peel, this compounds are effective against many strains of bacteria.

**Reference**

1. Bouchard, N.C.; Howland, M.A.; Greller, H.A.; Hoffman, R.S. and Nelson, L.S. (2005). Ischemic stroke associated with use of an ephedra-free dietary supplement containing synephrine. *Mayo Clin. Proc.*, 80, 541–545.
2. Haaz, S.; Fontaine, K.R.; Cutter, G.; Limdi, N.; Perumean-Chaney, S. and Allison, D.B. (2006). Citrus aurantium and synephrine alkaloids in the treatment of overweight and obesity: an update. *Obes. Rev.*, 7(1), 79–88.
3. Stohs, S.J.; Preuss, H.G. and Shara, M. (2011). The safety of Citrus aurantium (bitter orange) and its primary protoalkaloid p-synephrine. *Phytother. Res.*, 25, 1421– 1428.
4. Moraes, T.M.; Kushima, H.; Moleiro, F.C.; Santos, R.C. and Rocha, L.R. et al. (2009). Effects of limonene and essential oil from Citrus aurantium on gastric mucosa: Role of prostaglandins and gastric mucus secretion. *Chem-Bio. Interact.*, 180(3), 499-505.
5. Burt, S.A. (2004). Essential oils: Their antibacterial properties and potential applications in foods: A review. *Inter. J. Food Microbiol.*, 94(3), 223-253.
6. Ortuno, A.A.; Baidez, P.; Gomez, M.C.; Arcas, I.; Porras, A.G. and Del Rio, J.A. (2006). Citrus paradise and Citrus sinensis flavonoids: Their influence in the defence mechanism against *Penicillium digitatum*. *Food Chem.*, 98(2), 351-358.

**Phytochemical Analysis and Inhibitory Effect of *Citrus aurantium* L.****(Bitter Orange) Leaves on some Bacterial Isolates *in vitro*****Khudhair A. M. Abed AL Ani****Tareq R. Minnat****Osama K. Jalyl**

7. Akerle Heywood, O.V. and Synge, H. (1991). The conservation of medicinal plants. Cambridge.
8. Iniaghe, O.M.; Malomo, S.O. and Adebayo, J.O. (2009). Proximate composition and phytochemical constituents of leaves of some Acalypha species. Pak. J. Nutr, 8(3), 256–258.
9. Oloyede, O.I. (2005). Chemical profile of unripe pulp of *Carica papaya*. Pak. J. Nutr, 4(6), 379-381.
10. Mathew, B.; Jatawa, S.K. and Tiwaari A. (2012). Phytochemical analysis of *Citrus limonum* pulp and peel. International Journal of Pharmacy and Pharmaceutical Sciences, 4(2), 269-371.
11. Al-Juraifani, A.A. (2011). Antimicrobial activity of some medicinal plants used in Saudi Arabia. Can J Pure Appl Sci, 5(2), 1509-1512.
12. Cohen, M.L.(1992). Epidemiology of drug resistance: implications for a postantimicrobial era. Science, 257(5073), 1050-1055.
13. Chopra, R.N.; Nayar, S.L. and Chopra, I.C. (1956). Glossary of Indian medicinal plants. (Council of Scientific and Industrial Research, New Delhi).
14. Harborne, J.B. (1984). Phytochemical methods a guide to modern techniques of plant analysis. Chapman and Hill, London, p.5.
15. Jaffer, H.J.; Mahmud, M.J.; Jawad, A.M.; Naji, A. and Al-Naib, A. (1988). Phytochemical and biological screening of some Iraqi plants. Fitoterapia J.,59(3), 229-233.
16. Harborne, J.B. (1990). Phytochemical methods a guide to modern techniques of plant analysis .Chapman and Hill, London, 42(2), p115.
17. Al-Bid, M.R. (1985). Zurrzu samma msetzungder abschla B membrane in phoenix dactylifera. Wurzburg University. Wuzzburg F.R. of Germany.
18. Odebiyi, O.O. and Sofowora, E.A. (1978). Phytochemical screening of Nigerian medical plants II. Lloydia . Europe PMC plus, 41(3), 234–246.
19. Forbes, B.A. Sahm, D.F. and Weissfeld, A.S. (2007). Bailey and Scotts' Diagnostic microbiology 12th. ed., Elsevier China.

Phytochemical Analysis and Inhibitory Effect of *Citrus aurantium* L.  
(Bitter Orange) Leaves on some Bacterial Isolates *in vitro*

Khudhair A. M. Abed AL Ani

Tareq R. Minnat

Osama K. Jalyl

20. Bauer, A.W.; Kirby, W.M.M.; Sherris, J.C. and Turk, M. (1966). Antibiotic susceptibility testing by a standardized single disc method. *Am. J. Clin. Pathol*, 45(4), 493-496.
21. Srinivasan, D.; Nathan, S.; Suresh, T. and Lakshmana Perumalsamy, P. (2001). Antimicrobial activity of certain Indian Medicinal Plants used in folkloric Medicine. *J. Ethanopharmacol.*, 74(3), 217-220.
22. Masika, P.J. and Afolayan, A.J. (2002). Antimicrobial activity of some plants use for the treatment of livestock disease in the Eastern Cape. South Africa. *Journal of Ethanopharmacology*, 83, 129-134.
23. Karaman, I.; Sahin, F.; Güllüce, M.; Ögütçü, H.; Şengül, M. and Adıgüzel, A. (2003). Antimicrobial activity of aqueous and methanol extracts of *Juniperus Oxycedrus* L. *Journal of Ethanopharmacology*, 85, 231-235.
24. Steel, R.G. and Torrie, J.H. (1984). Principles and procedures of statistics a Biometrical Approach, 2nd ed., McGraw –Hill, Inc., Singapore, p.183.
25. Al-Badrani, B.A. (2002). Toxicological and pharmacological effects of Sibabhah (*Meliaazedarach*) and Oranamental Za-Rur (*Cotoneaster prostrate*) fruits. PhD thesis, College of Veterinary Medicine, University of Mosul.
26. Sofowra, A. (1993). Medicinal Plants and traditional Medicine in Africa. Spectrum Books Ltd., Ibadan, Nigeria. pp. 191-289.
27. Chede, P.S. (2013). Phytochemical analysis of *Citrus sinensis* peel. *International Journal of Pharma and Bio Sciences*, 4(1), 339 – 343.
28. Lawal, D.; Bala, J.A.; Aliyu, S.Y. and Huguma, M. A. (2013). Phytochemical screening and *in vitro* anti-bacterial studies of the ethanolic extract of *Citrus sinensis* (Linn.) peel against some clinical bacterial isolates. *International Journal of Innovation and Applied Studies*, 2(2), 138 -145.
29. Swapnil, M.; Rakesh, V.; Bhupendra, V.; Tusharbindu, D.; Vishal, P. and Devang, P. (2011). Pharmacognostic and Phytochemical Characterization of Leaves of *Citrus maxima*. *International Journal of Innovative Pharmaceutical Research*, 2(4), 175-178.

Phytochemical Analysis and Inhibitory Effect of *Citrus aurantium* L.  
(Bitter Orange) Leaves on some Bacterial Isolates *in vitro*

Khudhair A. M. Abed AL Ani

Tareq R. Minnat

Osama K. Jalyl

30. Ahmad, M.M.; Salim-Ur-Rehman, Zafar, I.; Faqir, M. A. and Javaid, I. (2006). Genetic variability to essential oil composition in four Citrus fruit species. *Pakistan Journal of Botany*, 38(2), 319-324.
31. Kirbaşlar, G.F.; Tavman, A.; Dülger, B. and Türker, G. (2009). Antimicrobial activity of Turkish citrus peel oils. *Pakistan Journal of Botany*, 41(6), 3207-3212.
32. Bouabdelli, F.; Djelloul, A.; Kaid-Omar, Z.; Semmoud, A. and Addou, A. (2012). Antimicrobial Activity of 22 Plants Used in Urolithiasis Medicine in Western Algeria. *Asian Pacific Journal of Tropical Disease*, 12(1), 530-535.
33. Gopal, P.V. (2012). Evaluation of antimicrobial activity of *Citrus aurantium* against some gram positive and negative bacterial strains. *Pharmacia*, 1(3), 107-109.
34. Kabra, A.O.; Bairagi, G.B.; Mahamuni, A.S. and Wanare, R.S. (2012). In vitro antimicrobial activity and phytochemical analysis of the peels of *Citrus medica* L. *International Journal of Research in Pharmaceutical and Biomedical Sciences*, 3 (1), 34-37.
35. Cano, A.; Medinaan, A. and Bermejo, A. (2008). Bioactive compounds in different citrus varieties. Discrimination among cultivars. *Journal of Food Composition and Analysis*, 21(5), 377- 381.
36. Dhanavade, M.J.; Chidamber, B.; Jalkute, J.; Ghosh, S. and Sonawane, K.D. (2011). Study Antimicrobial Activity of Lemon (*Citrus lemon* L.) Peel Extract. *British Journal of Pharmacology and Toxicology*, 2(3), 119-122.