

Retarded Integral Inequalities with Iterated Integrals

Ali W.K. Sangawi<sup>1,2</sup> and Sudad M. Rasheed<sup>1,2</sup>

Retarded Integral Inequalities with Iterated Integrals

Ali W.K. Sangawi<sup>1,2</sup> and Sudad M. Rasheed<sup>1,2</sup>

Received 9 October 2013 ; Accepted 21 October 2014

**Abstract**

This paper presents an iterated and iterated retarded integral inequalities and explicit bounds to unknown functions on some iterated and iterated retarded integral inequalities are established.

**keywords:** Integral inequalities, Retarded integral inequalities, Non-decreasing functions, Non-negative continuous functions, partial derivatives, Explicit Bounds.

**تكامل المترجمات المتكررة والتباطؤية**

د. علي وهاب كريم – قسم الرياضيات – كلية العلوم – جامعة السليمانية

E-mail : [alisangawi2000@yahoo.com](mailto:alisangawi2000@yahoo.com)

م. سوّدد موسى رشيد – قسم الرياضيات – كلية العلوم – جامعة السليمانية

E-mail : [sudadmusa@hotmail.com](mailto:sudadmusa@hotmail.com)

**المخلص**

ان الهدف الرئيسي من هذا البحث هو تقديم المترجمات التكاملية المتكررة واعطاء قيود صريحة للدوال المجهولة في بعض المترجمات التكاملية المتكررة والمترجمات التكاملية المتكررة التباطؤية .

### Introduction

Integral inequalities with iterated integrals are indispensable for us in the quantitative study of various differential equations and integral equations. motivated by a desire to apply integral inequalities which provide explicit bounds on unknown functions, in the development of the theory of differential and integral equations with retarded arguments.

#### **Lemma 1 .**

Let  $u(x)$  and  $g(x)$  be nonnegative continuous functions on  $I = [0, \infty)$  for the inequality

$$u(x) \leq c + \int_a^x g(t)u(t)dt \quad , \quad t \in I,$$

holds, where  $c$  is constant. Then

$$u(x) \leq c e^{\int_a^x g(t)dt} \quad , \quad t \in I.$$

The result was proved by Gronwall [13]. Gronwall type integral inequalities provide a necessary tool for the study of the theory of differential equations, integral equations and inequalities of various types see [2-11, 15 ].Some Gronwall-Bellman type integral inequalities with fixed delay has been presented in [1].

The aim of the present paper is to establish explicit bounds on more general integral inequalities with iterated and iterated retarded integral inequalities.

The plan of the paper is as follows: Section 2 presents some iterated integral inequalities. Section 3 presents some iterated retarded integral inequalities. Finally, Section 4 presents a short conclusion.

### Integral Some Iterated Inequalities

This section presents some iterated integral inequalities and then give explicit bounds to unknown functions. Later on  $\mathbb{R}_+ = [0, \infty)$ ,  $I = [\alpha, \beta]$  and  $I_t = [t, \beta]$ .

Retarded Integral Inequalities with Iterated Integrals

Ali W.K. Sangawi<sup>1,2</sup> and Sudad M. Rasheed<sup>1,2</sup>

**Theorem : (2.1)**

Let

$u(t), f(t), a(t), g(t), h(t) \in C(\mathbb{R}_+, \mathbb{R}_+), k(t, s), k_r(t, s) \in C(D, \mathbb{R}_+)$  and  $p > 1$

be real constant, where

$$D = \{(t, s) \in \mathbb{R}_+^2; 0 \leq s \leq t \leq \infty\}.$$

**(a<sub>1</sub>)**

If  $u^p(t) \leq a(t) + \int_0^t f(s) [u^p(s) + \int_0^s k(s, \sigma)u(\sigma)d\sigma] ds$  , .....(2.1)

for  $t \in \mathbb{R}_+$  , then

$$u(t) \leq \left[ a(t) + \int_0^t A_1(s) \exp^{\int_s^t B_1(\tau) d\tau} ds \right]^{\frac{1}{p}} , \dots\dots\dots (2.2)$$

for  $t \in \mathbb{R}_+$  , where  $A_1(t) = \frac{f(t)}{p} [pa(t) + \int_0^t k(t, \sigma) [(p-1) + a(\sigma)] d\sigma]$  and

$$B_1(t) = f(t) \left[ 1 + \int_0^t \frac{k(t, \sigma)}{p} d\sigma \right].$$

**(a<sub>2</sub>)**

Let  $c(t)$  be real-valued positive continuous and nondecreasing function defined in  $\mathbb{R}_+$ .

If  $u^p(t) \leq c^p(t) + \int_0^t f(s) \{u^{(p)}(s) + \int_0^s k(s, \sigma)u(\sigma)d\sigma\} ds$  , ..... (2.3)

for  $t \in \mathbb{R}_+$  , then

$$u(t) \leq c(t) \left[ 1 + \int_0^t A_2(s) \exp^{\int_s^t B_2(\tau) d\tau} ds \right]^{\frac{1}{p}} \dots\dots\dots (2.4)$$

for  $t \in \mathbb{R}_+$  , where

$$A_2(t) = f(t) \left[ 1 + \int_0^t k(t, \sigma) c^{1-p}(\sigma) d\sigma \right]$$

and

$$B_2(t) = f(t) \left[ 1 + \int_0^t \frac{\{k(t, \sigma)c^{1-p}(\sigma)\}}{p} d\sigma \right].$$

**(a<sub>3</sub>)**

Retarded Integral Inequalities with Iterated Integrals

Ali W.K. Sangawi<sup>1,2</sup> and Sudad M. Rasheed<sup>1,2</sup>

If

$$u^p(t) \leq a(t) + \int_0^t k(t,s) \left\{ u(s) + \int_0^s [g(\sigma)u^p(\sigma) + h(\sigma)u(\sigma)] d\sigma \right\} ds ,$$

..... (2.5)

for  $t \in \mathbb{R}_+$ , then

$$u(t) \leq \left[ a(t) + \int_0^t A_3(s) \exp \int_s^t B_3(\tau) d\tau ds \right]^{\frac{1}{p}} \dots\dots (2.6)$$

for  $t \in \mathbb{R}_+$ , where

$$\begin{aligned} A_3(t) = & \frac{k(t,t)}{p} \left[ (p-1) + a(t) + \int_0^t [pg(\sigma)a(\sigma) + h(\sigma)[(p-1) + a(\sigma)]] d\sigma \right] \\ & + \int_0^t \frac{k_t(t,s)}{p} \left[ (p-1) + a(s) \right. \\ & + \left. \int_0^s [pg(\sigma)a(\sigma) + h(\sigma)[(p-1) + a(\sigma)]] d\sigma \right] ds, \end{aligned}$$

And

$$B_3(t) = \frac{k(t,t)}{p} \left[ 1 + \int_0^t [pg(\sigma) + h(\sigma)] d\sigma \right] + \int_0^t \frac{k_t(t,s)}{p} \left[ 1 + \int_0^s [pg(\sigma) + h(\sigma)] d\sigma \right] ds.$$

**Proof:**

(a<sub>1</sub>)

Defin a function  $z(t)$  by  $z(t) = \int_0^t f(s) \left\{ u^p(s) + \int_0^s k(s,\sigma)u(\sigma) d\sigma \right\} ds.$

..... (2.7)

Then  $z(t) \geq 0$ ,  $z(t)$  is nondecreasing for  $t \in I$  and inequality (2.1) can be written as

Retarded Integral Inequalities with Iterated Integrals

Ali W.K. Sangawi<sup>1,2</sup> and Sudad M. Rasheed<sup>1,2</sup>

$$u^p(t) \leq a(t) + z(t). \quad \dots (2.8)$$

From inequality (2.8) and using the elementary inequality see [14], [12]

$$x^{\frac{1}{p}}y^{\frac{1}{q}} \leq \frac{x}{p} + \frac{y}{q}$$

Where  $x, y \geq 0$  and  $\frac{1}{p} + \frac{1}{q} = 1$ ,

we observe that

$$u(t) \leq [a(t) + z(t)]^{\frac{1}{p}} (1)^{\frac{\frac{1}{p}}{p-1}}$$

$$\leq \frac{(p-1)}{p} + \frac{a(t)}{p} + \frac{z(t)}{p}, \quad \dots (2.9)$$

Differentiating (2.7) and using (2.8), (2.9) we get:

$$z'(t) = f(t) \left\{ u^p(t) + \int_0^t k(t, \sigma) u(\sigma) d\sigma \right\}$$

$$\leq f(t) \left\{ a(t) + z(t) + \int_0^t k(t, \sigma) \left[ \frac{(p-1)}{p} + \frac{a(\sigma)}{p} + \frac{z(\sigma)}{p} \right] d\sigma \right\}$$

$$= \frac{f(t)}{p} \left\{ pa(t) + \int_0^t k(t, \sigma) [(p-1) + a(\sigma)] d\sigma \right\}$$

$$+ f(t) \left\{ 1 + \int_0^t \frac{k(t, \sigma)}{p} d\sigma \right\} z(t)$$

$$= A_1(t) + B_1(t)z(t).$$

Integrating both sides of the above inequality from 0 to t we get :

$$z(t) \leq \int_0^t A_1(s) \exp^{-\int_s^t B_1(\tau) d\tau} ds \quad \dots (2.10)$$

Using (2.10) in (2.8), we get the required inequality in (2.2).

## Retarded Integral Inequalities with Iterated Integrals

Ali W.K. Sangawi<sup>1,2</sup> and Sudad M. Rasheed<sup>1,2</sup>**(a<sub>2</sub>)**

Since  $c(t)$  is positive, continuous and nondecreasing function for  $t \in \mathbb{R}_+$ , from (2.3) then one can get :

$$\left(\frac{u(t)}{c(t)}\right)^p \leq 1 + \int_0^t f(s) \left\{ \left(\frac{u(s)}{c(s)}\right)^p + \int_0^s k(s,\sigma) c^{1-p}(\sigma) \frac{u(\sigma)}{c(\sigma)} d\sigma \right\} ds$$

Now an application of the inequality given in (a<sub>1</sub>) yields the desired result in (2.4).

**(a<sub>3</sub>)**

Define the function  $z(t)$  by

$$z(t) = \int_0^t k(t,s) \left\{ u(s) + \int_0^s [g(\sigma)u^p(\sigma) + h(\sigma)u(\sigma)] d\sigma \right\} ds$$

....(2.11)

Then as in the proof of part (a<sub>1</sub>), from (2.11) we see that the inequalities (2.8) and (2.9) hold.

Differentiating (2.11) and using (2.8), (2.9) and the fact that  $z(t)$  is nondecreasing in  $t$  we get:

$$z'(t) = k(t,t) \left\{ u(t) + \int_0^t [g(\sigma)u^p(\sigma) + h(\sigma)u(\sigma)] d\sigma \right\} + \int_0^t k_t(t,s) \left\{ u(s) + \int_0^s [g(\sigma)u^p(\sigma) + h(\sigma)u(\sigma)] d\sigma \right\} ds$$

$$\leq k(t,t) \left\{ \frac{p-1}{p} + \frac{a(t)}{p} + \frac{z(t)}{p} + \int_0^t \left[ g(\sigma)[a(\sigma) + z(\sigma)] + h(\sigma) \left[ \frac{p-1}{p} + \frac{a(\sigma)}{p} + \frac{z(\sigma)}{p} \right] \right] d\sigma \right\}$$

Retarded Integral Inequalities with Iterated Integrals

Ali W.K. Sangawi<sup>1,2</sup> and Sudad M. Rasheed<sup>1,2</sup>

$$\begin{aligned}
 & + \int_0^t k_t(t,s) \left\{ \frac{p-1}{p} + \frac{a(s)}{p} + \frac{z(s)}{p} \right. \\
 & \quad \left. + \int_0^s [g(\sigma)[a(\sigma) + z(\sigma)] + h(\sigma) \left[ \frac{p-1}{p} + \frac{a(\sigma)}{p} \right. \right. \\
 & \quad \left. \left. + \frac{z(\sigma)}{p} \right] d\sigma \right\} ds \\
 & = \frac{k(t,t)}{p} \left\{ (p-1) + a(t) \right. \\
 & \quad \left. + \int_0^t [pg(\sigma)a(\sigma) + h(\sigma)[(p-1) + a(\sigma)]] d\sigma \right\} \\
 & + \int_0^t \frac{k_t(t,s)}{p} \left\{ (p-1) + a(s) \right. \\
 & \quad \left. + \int_0^s [pg(\sigma)a(\sigma) \right. \\
 & \quad \left. + h(\sigma)[(p-1) + a(\sigma)]] d\sigma \right\} ds + \frac{k(t,t)}{p} \\
 & \left\{ z(t) + \int_0^t [pg(\sigma)z(\sigma) + h(\sigma)z(\sigma)] d\sigma \right\} \\
 & \quad + \int_0^t \frac{k_t(t,s)}{p} \left\{ z(s) \right. \\
 & \quad \left. + \int_0^s [pg(\sigma)z(\sigma) + h(\sigma)z(\sigma)] d\sigma \right\} ds \\
 & = A_3(t) + B_3(t)z(t).
 \end{aligned}$$

Integrating both sides of the above inequality from 0 to  $t$  yields

$$z(t) \leq \int_0^t A_3(s) \exp \int_s^t B_3(\tau) d\tau ds . \quad \dots (2.12)$$

Retarded Integral Inequalities with Iterated Integrals

Ali W.K. Sangawi<sup>1,2</sup> and Sudad M. Rasheed<sup>1,2</sup>

Using (2.12) in (2.8), we get the required inequality in (2.6).

**Theorem : (2.2)**

Let  $u(t), g(t) \in C(I, \mathbb{R}_+)$ ,  $k(t, s), b(t, s), c(t, s) \in C(D, \mathbb{R}_+)$ ,

$h(t, s, \sigma) \in C(E, \mathbb{R}_+)$  and  $a(t), a'(t) \in C(I, \mathbb{R}_+)$ ,  $p > 1$  be real constant.

(b<sub>1</sub>)

Let  $\phi(t) \in C(I, \mathbb{R}_+)$  and  $u^p(t) \leq a(t) + \int_{\alpha}^t \phi(s)u(s)ds + \int_{\alpha}^t \int_{\alpha}^s k(s, \tau)u(\tau)d\tau ds +$

$$\int_{\alpha}^t \int_{\alpha}^s \int_{\alpha}^{\tau} h(s, \tau, \sigma)u(\sigma)d\sigma d\tau ds + \int_{\alpha}^t \int_{\alpha}^s c(s, \tau)u(\tau)d\tau ds \dots (2.13)$$

for  $t \in I$ . If

$$P_1 = \frac{1}{p} \int_{\alpha}^{\beta} \int_{\alpha}^s c(s, \tau) \exp^{\int_{\alpha}^{\tau} B_4(\xi)d\xi} d\tau ds < 1 \dots (2.14)$$

then

$$u(t) \leq \left[ a(t) + M_1 \exp^{\int_{\alpha}^t B_4(\xi)d\xi} + \int_{\alpha}^t A_2(\eta) \exp^{\int_{\alpha}^{\eta} B_4(\xi)d\xi} d\eta \right]^{\frac{1}{p}} \dots (2.15)$$

for  $t \in I$ , where

$$A_2(t) = \frac{1}{p} \{ \phi(t)[(p-1) + a(t)] + \int_{\alpha}^t k(t, \tau) [(p-1) + a(\tau)] d\tau + \int_{\alpha}^t \int_{\alpha}^{\tau} h(t, \tau, \sigma) [(p-1) + a(\sigma)] d\sigma d\tau \},$$

$$B_4(t) = \frac{1}{p} \left\{ \phi(t) \mid \int_{\alpha}^t k(t, \tau) d\tau \mid \int_{\alpha}^t \int_{\alpha}^{\tau} h(t, \tau, \sigma) d\sigma d\tau \right\}$$

and

$M_1 =$



Retarded Integral Inequalities with Iterated Integrals

Ali W.K. Sangawi<sup>1,2</sup> and Sudad M. Rasheed<sup>1,2</sup>

$$\frac{1}{1 - P_1} \left\{ \frac{1}{p} \int_{\alpha}^{\beta} \int_{\alpha}^{\xi} c(s, \tau) [(p - 1) + a(\tau) + \int_{\alpha}^{\tau} A_4(\eta) \exp^{\int_{\alpha}^{\xi} B_4(\xi) d\xi} d\eta] d\tau ds \right\}$$

(b<sub>2</sub>)

Let  $k(t, s)$ ,  $b(t, s)$ ,  $h_2(t, s, \sigma)$  are nondecreasing in  $t \in I$ , for each  $s \in I$  and

$$u^p(t) \leq a(t) + \int_{\alpha}^t k(t, \tau) u(\tau) d\tau + \int_{\alpha}^t \int_{\alpha}^{\xi} h(t, s, \sigma) u(\sigma) d\sigma ds + \int_{\alpha}^{\beta} b(t, s) \int_{\alpha}^{\xi} c(s, \tau) u(\tau) d\tau ds$$

for  $t \in I$ . ..... (2.16)

If  $P_2 = \frac{1}{p} \int_{\alpha}^{\beta} b(t, s) \int_{\alpha}^{\xi} c(s, \tau) \exp^{\int_{\alpha}^{\xi} B_2(\eta, \tau) d\eta} d\tau ds < 1$  ..... (2.17)

then

$$u(t) \leq \left[ a(t) + M_2 \exp^{\int_{\alpha}^t B_2(\eta, t) d\eta} + \int_{\alpha}^t A_5(\xi, t) \exp^{\int_{\alpha}^{\xi} B_2(\eta, t) d\eta} d\xi \right]^{\frac{1}{p}}$$

..... (2.19)

for  $t \in I$ , where

$$A_5(t, T) = \frac{1}{p} \left\{ k(T, t) [(p - 1) + a(t)] + \int_{\alpha}^t h(T, t, \sigma) [(p - 1) + a(\sigma)] d\sigma \right\}$$

$$B_5(t, T) = \frac{1}{p} \left\{ k(T, t) + \int_{\alpha}^t h(T, t, \sigma) d\sigma \right\}$$

and

$$M_2 =$$

Retarded Integral Inequalities with Iterated Integrals

Ali W.K. Sangawi<sup>1,2</sup> and Sudad M. Rasheed<sup>1,2</sup>

$$\frac{1}{1-p_2} \left\{ \frac{1}{p} \int_{\alpha}^{\beta} b(t, s) \int_{\alpha}^{\sigma} c(s, \tau) [(p-1) + a(\tau) + \int_{\alpha}^{\tau} A_3(\xi, \tau) \exp^{\int_{\xi}^{\tau} B_3(\eta, \tau) d\eta} d\xi] d\tau ds \right\}$$

(b<sub>3</sub>)

Let  $r(t) \in C(I, \mathbb{R}_+)$  and

$$u^p(t) \leq a(t) + \int_{\alpha}^t g(s) \left\{ u(s) + \int_{\sigma}^s k(s, \sigma) u(\sigma) d\sigma + \int_{\alpha}^{\beta} r(\sigma) u(\sigma) d\sigma \right\} \dots\dots\dots(2.20)$$

for  $t \in I$ . If

$$P_3 = \int_{\alpha}^{\beta} r(\sigma) \exp^{\int_{\alpha}^{\sigma} B_3(\eta) d\eta} d\sigma < 1, \dots\dots (2.21)$$

then

$$u(t) \leq \left[ a(t) + M_3 \exp^{\int_{\alpha}^t B_3(\eta) d\eta} + \int_{\alpha}^t A_3(\xi) \exp^{\int_{\alpha}^{\xi} B_3(\eta) d\eta} d\xi \right]^{\frac{1}{p}}, \dots (2.21)$$

for  $t \in I$ , where

$$A_3(t) = g(t) \frac{1}{p} \left\{ (p-1) + a(t) + \int_{\alpha}^t k(t, \sigma) [(p-1) + a(\sigma)] d\sigma + \int_{\alpha}^{\beta} r(\sigma) [(p-1) + a(\sigma)] d\sigma \right\}$$

$$B_3(t) = \frac{1}{p} g(t) + k(t, t) + \int_{\alpha}^t k_{\tau}(t, \sigma) d\sigma \quad \text{and}$$

$$M_3 = \frac{1}{1-P_3} \left\{ \int_{\alpha}^{\beta} r(\sigma) \int_{\alpha}^{\sigma} A_3(\xi) \exp^{\int_{\xi}^{\sigma} B_3(\eta) d\eta} d\xi d\sigma \right\}.$$

**Proof :**

(b<sub>1</sub>)

Define a function  $z(t)$  by

Retarded Integral Inequalities with Iterated Integrals

Ali W.K. Sangawi<sup>1,2</sup> and Sudad M. Rasheed<sup>1,2</sup>

$$\begin{aligned}
 z(t) = & \int_{\alpha}^t \phi(s)u(s)ds + \int_{\alpha}^t \int_{\alpha}^s k(s, \tau)u(\tau)d\tau ds \\
 & + \int_{\alpha}^t \int_{\alpha}^s \int_{\alpha}^{\tau} h(s, \tau, \sigma)u(\sigma)d\sigma d\tau ds \\
 & + \int_{\alpha}^{\beta} \int_{\alpha}^s c(s, \tau)u(\tau)d\tau ds
 \end{aligned}
 \dots\dots\dots (2.22)$$

Then  $z(t) \geq 0$  ,  $z(t)$  is nondecreasing for  $t \in I$

$$z(\alpha) = \int_{\alpha}^{\beta} \int_{\alpha}^s c(s, \tau)u(\tau)d\tau ds \dots\dots\dots(2.23)$$

Then as in the proof of part (a<sub>1</sub>), from (2.23)we see that the inequalities(2.8)and(2.9)hold.

Differentiating(2.23)and using(2.9)and the fact that  $z(t)$  is nondecreasing in  $t$ , we get:

$$\begin{aligned}
 z'(t) = & \phi(t)u(t) + \int_{\alpha}^t k(t, \tau)u(\tau)d\tau \\
 & + \int_{\alpha}^t \int_{\alpha}^{\tau} h(t, \tau, \sigma)u(\sigma)d\sigma d\tau \\
 \leq & \phi(t)\frac{1}{p}[(p-1) + a(t) + z(t)] + \int_{\alpha}^t k(t, \tau)\frac{1}{p}[(p-1) \\
 & + a(\tau) + z(\tau)] d\tau \\
 & + \int_{\alpha}^t \int_{\alpha}^{\tau} h(t, \tau, \sigma)\frac{1}{p}[(p-1) + a(\sigma) + z(\sigma)]d\sigma d\tau \\
 = & \frac{1}{p}\{ \phi(t)[(p-1) + a(t)] + \int_{\alpha}^t k(t, \tau) [(p-1) + a(\tau)]d\tau \\
 & + \int_{\alpha}^t \int_{\alpha}^{\tau} h(t, \tau, \sigma) [(p-1) + a(\sigma)]d\sigma d\tau \}
 \end{aligned}$$

Retarded Integral Inequalities with Iterated Integrals

Ali W.K. Sangawi<sup>1,2</sup> and Sudad M. Rasheed<sup>1,2</sup>

$$\begin{aligned}
 & + \frac{1}{p} \left\{ \phi(t)z(t) + \int_{\alpha}^t k(t, \tau)z(\tau)d\tau \right. \\
 & \left. + \int_{\alpha}^t \int_{\alpha}^{\tau} h(t, \tau, \sigma)z(\sigma)d\sigma d\tau \right\}.
 \end{aligned}$$

But  $z(t)$  is nonnegative and nondecreasing for  $t \in I$ , then

$$z'(t) \leq A_4(t) + B_4(t)z(t)$$

Therefore,  $\alpha \leq \eta \leq t \leq \beta$ , one can have :

$$\frac{d}{d\eta} \left[ z(\eta) \exp^{\int_{\eta}^t B_4(\xi) d\xi} \right] \leq A_4(\eta) \exp^{\int_{\eta}^t B_4(\xi) d\xi}$$

Integrating both sides of the above inequality from  $\alpha$  to  $t$ , for  $t \in I$ , we get:

$$z(t) \leq z(\alpha) \exp^{\int_{\alpha}^t B_4(\xi) d\xi} + \int_{\alpha}^t A_4(\eta) \exp^{\int_{\eta}^t B_4(\xi) d\xi} d\eta, \dots\dots(2.24)$$

from (2.24) and (2.9) one can get

$$\begin{aligned}
 u(t) & \leq \frac{1}{r} z(\alpha) \exp^{\int_{\alpha}^t B_4(\xi) d\xi} + \frac{1}{r} [(p-1) + a(t) + \\
 & \int_{\alpha}^t A_4(\eta) \exp^{\int_{\eta}^t B_4(\xi) d\xi} d\eta] \\
 & \dots\dots(2.25)
 \end{aligned}$$

From (2.23) and (2.25) Which implies

$$\begin{aligned}
 z(\alpha) & \leq \int_{\alpha}^{\beta} \int_{\alpha}^s c(s, \tau) \left\{ \frac{1}{p} z(\alpha) \exp^{\int_{\alpha}^t B_4(\xi) d\xi} \right. \\
 & \left. + \frac{1}{p} [(p-1) + a(\tau) \right. \\
 & \left. + \int_{\alpha}^{\tau} A_4(\eta) \exp^{\int_{\eta}^t B_4(\xi) d\xi} d\eta] \right\} d\tau ds
 \end{aligned}$$

Retarded Integral Inequalities with Iterated Integrals

Ali W.K. Sangawi<sup>1,2</sup> and Sudad M. Rasheed<sup>1,2</sup>

then

$$z(\alpha) \left[ 1 - \frac{1}{p} \int_{\alpha}^{\beta} \int_{\alpha}^s c(s, \tau) \exp^{\int_{\alpha}^{\tau} B_4(\xi) d\xi} d\tau ds \right] \leq \int_{\alpha}^{\beta} \int_{\alpha}^s c(s, \tau) \frac{1}{p} [(p-1) + \alpha(\tau) + \int_{\alpha}^{\tau} A_4(\eta) \exp^{\int_{\alpha}^{\eta} B_4(\xi) d\xi} d\eta] d\tau ds$$

from (2.14) we obtain that

$$z(\alpha) \leq M_1. \dots\dots\dots(2.26)$$

The required inequality (2.15) follows from (2.26), (2.24) and (2.8).

(b<sub>2</sub>)

Fix any  $T, \alpha \leq T \leq \beta$ , then for  $\alpha \leq t \leq T$ , we have

$$u^p(t) \leq a(t) + \int_{\alpha}^t k(T, \tau) u(\tau) d\tau + \int_{\alpha}^t \int_{\alpha}^s h(T, s, \sigma) u(\sigma) d\sigma ds + \int_{\alpha}^{\beta} b(T, s) \int_{\alpha}^s c(s, \tau) u(\tau) d\tau ds, \dots\dots\dots(2.27)$$

Define a Function  $z(t, T), \alpha \leq t \leq T$  by

$$z(t, T) = \int_{\alpha}^t k(T, \tau) u(\tau) d\tau + \int_{\alpha}^t \int_{\alpha}^s h(T, s, \sigma) u(\sigma) d\sigma ds + \int_{\alpha}^{\beta} b(T, s) \int_{\alpha}^s c(s, \tau) u(\tau) d\tau ds \dots\dots\dots(2.28)$$

Then  $z(t, T) \geq 0, z(t, T)$  is nondecreasing for  $t \in I$ ,

$$z(\alpha, T) = \int_{\alpha}^{\beta} b(T, s) \int_{\alpha}^s c(s, \tau) u(\tau) d\tau ds, \dots\dots\dots(2.29)$$

and inequality (2.16) can be written as

$$u^p(t) \leq a(t) + z(t, T), \quad \alpha \leq t \leq T. \dots\dots\dots(2.30)$$

Retarded Integral Inequalities with Iterated Integrals

Ali W.K. Sangawi<sup>1,2</sup> and Sudad M. Rasheed<sup>1,2</sup>

Then as in the proof of part (a<sub>1</sub>), from (2.30) we see that the inequalities (2.31) hold.

$$u(t) \leq \frac{(p-1)}{p} + \frac{a(t)}{p} + \frac{z(t,T)}{p}, \quad \alpha \leq t \leq T$$

..... (2.31)

Differentiating (2.28) and using (2.31) and the fact that  $z(t, T)$  is nondecreasing in  $t$ , we get

$$\begin{aligned} z'(t, T) - k(T, t)u(t) &+ \int_{\alpha}^t h(T, t, \sigma)u(\sigma) d\sigma \\ &\leq \frac{1}{p} \{ k(T, t)[(p-1) + a(t)] \\ &+ \int_{\alpha}^t h(T, t, \sigma) [(p-1) + a(\sigma)] d\sigma \} + \frac{1}{p} \{ k(T, t) \\ &+ \int_{\alpha}^t h(T, t, \sigma) d\sigma \} z(t, T), \end{aligned}$$

then

$$z'(t, T) \leq A_{\bar{z}}(t, T) + B_{\bar{z}}(t, T)z(t, T) \quad \text{.....(2.32)}$$

for  $\alpha \leq T$  by setting  $t = \eta$  in (2.32) and integrating it with respect to  $\eta$  from  $\alpha$  to  $T$ , we get:

$$\begin{aligned} z(T, T) &\leq \\ z(\alpha, T) \exp^{\int_{\alpha}^T B_{\bar{z}}(\xi, T) d\xi} &+ \int_{\alpha}^T A_{\bar{z}}(\eta, T) \exp^{\int_{\eta}^T B_{\bar{z}}(\xi, T) d\xi} d\eta, \end{aligned}$$

.....(2.33)

Since  $T$  is arbitrary from (2.33), (2.31), (2.30) and (2.29) with  $T$  replaced by  $t$  one can get

Retarded Integral Inequalities with Iterated Integrals

Ali W.K. Sangawi<sup>1,2</sup> and Sudad M. Rasheed<sup>1,2</sup>

$$z(t, t) \leq z(\alpha, t) \exp^{\int_{\alpha}^t B_{\beta}(\xi, t) d\xi} + \int_{\alpha}^t A_{\beta}(\eta, t) \exp^{\int_{\alpha}^t B_{\beta}(\xi, t) d\xi} d\eta, \dots\dots\dots(2.34)$$

$$u^p(t) \leq a(t) + z(t, t), \dots\dots\dots(2.35)$$

$$u(t) \leq \frac{1}{p} \{ (p-1) + a(t) + z(t, t) \}$$

$$\leq \frac{1}{p} \{ (p-1) + a(t) + z(\alpha, t) \exp^{\int_{\alpha}^t B_{\beta}(\xi, t) d\xi} + \int_{\alpha}^t A_{\beta}(\eta, t) \exp^{\int_{\alpha}^t B_{\beta}(\xi, t) d\xi} d\eta, \dots\dots\dots(2.36)$$

Where

$$z(\alpha, t) = \int_{\alpha}^{\beta} b(t, s) \int_{\alpha}^s c(s, \tau) u(\tau) d\tau ds \dots\dots\dots (2.37)$$

then from (2.37) and (2.36) Which implies

$$z(\alpha, t) \leq \int_{\alpha}^{\beta} b(t, s) \int_{\alpha}^s c(s, \tau) \frac{1}{p} \{ (p-1) + a(\tau) + z(\alpha, \tau) \exp^{\int_{\alpha}^{\tau} B_{\beta}(\xi, \tau) d\xi} + \int_{\alpha}^{\tau} A_{\beta}(\eta, \tau) \exp^{\int_{\alpha}^{\tau} B_{\beta}(\xi, \tau) d\xi} d\eta \} d\tau ds.$$

But  $z(t, T)$  is nonnegative and nondecreasing for  $t \in I$ , then

Retarded Integral Inequalities with Iterated Integrals

Ali W.K. Sangawi<sup>1,2</sup> and Sudad M. Rasheed<sup>1,2</sup>

$$z(\alpha, t) \left[ 1 - \frac{1}{p} \int_{\alpha}^{\beta} b(t, s) \int_{\alpha}^s c(s, \tau) \exp^{\int_{\alpha}^{\tau} B_s(\xi, \tau) d\xi} d\tau ds \right] \leq$$

$$\int_{\alpha}^{\beta} b(t, s) \int_{\alpha}^s c(s, \tau) \frac{1}{p} \left\{ (p-1) + \alpha(\tau) + \int_{\alpha}^{\tau} A_s(\eta, \tau) \exp^{\int_{\eta}^{\tau} B_s(\xi, \tau) d\xi} d\eta \right\} d\tau ds$$

from (2.17) we obtain that

$$z(\alpha, t) \leq M_2 . \dots\dots\dots(2.38)$$

The required inequality (2.18) follows from (2.38), (2.34) and (2.35).

(b<sub>3</sub>)

Define the function  $z(t)$  by

$$z(t) = \int_{\alpha}^t g(s) \left\{ u(s) + \int_{\alpha}^s k(s, \sigma) u(\sigma) d\sigma + \int_{\alpha}^{\beta} r(\sigma) u(\sigma) d\sigma \right\} ds . \dots\dots\dots(2.39)$$

Then  $z(t) \geq 0$ ,  $z(t)$  is non-decreasing for  $t \in I, z(\alpha) = 0$ .

Then as in the proof of part (a<sub>1</sub>). From (2.39) we see that the inequalities (2.8) and (2.9) hold. Differentiating (2.39) and using (2.9) and the fact that  $z(t)$  is nondecreasing in  $t$ , we get

$$z'(t) = g(t) \left[ u(t) + \int_{\alpha}^t k(t, \sigma) u(\sigma) d\sigma + \int_{\alpha}^{\beta} r(\sigma) u(\sigma) d\sigma \right]$$



Retarded Integral Inequalities with Iterated Integrals

Ali W.K. Sangawi<sup>1,2</sup> and Sudad M. Rasheed<sup>1,2</sup>

$$\begin{aligned} &\leq \frac{1}{p} g(t) \left\{ (p-1) + a(t) + \int_{\alpha}^t k(t, \sigma) [(p-1) + a(\sigma)] d\sigma \right. \\ &\quad \left. + \int_{\alpha}^{\beta} r(\sigma) [(p-1) + a(\sigma)] d\sigma \right\} \\ &+ \frac{1}{p} g(t) \left\{ z(t) + \int_{\alpha}^t k(t, \sigma) z(\sigma) d\sigma + \int_{\alpha}^{\beta} r(\sigma) z(\sigma) d\sigma \right\} \\ &= \\ &A_6(t) + \frac{1}{p} g(t) \left\{ z(t) + \int_{\alpha}^t k(t, \sigma) z(\sigma) d\sigma + \int_{\alpha}^{\beta} r(\sigma) z(\sigma) d\sigma \right\} \end{aligned}$$

Let

$$v(t) = z(t) + \int_{\alpha}^t k(t, \sigma) z(\sigma) d\sigma + \int_{\alpha}^{\beta} r(\sigma) z(\sigma) d\sigma \quad \dots\dots\dots (2.40)$$

then  $v(t) \geq 0$  and nondecreasing for  $t \in I$ , and since  $z(\alpha) = 0$  then

$$v(\alpha) = \int_{\alpha}^{\beta} r(\sigma) z(\sigma) d\sigma \quad \dots\dots\dots (2.41)$$

$$z(t) \leq v(t) \quad \dots\dots\dots (2.42)$$

$$z'(t) \leq A_6(t) + \frac{1}{p} g(t) v(t). \quad \dots\dots\dots (2.43)$$

Differentiating both sides of (2.40) and using (2.42) and (2.43), We get:

$$\begin{aligned} v'(t) &= z'(t) + k(t, t)z(t) + \int_{\alpha}^t k_t(t, \sigma) z(\sigma) d\sigma \\ &\leq A_6(t) + \frac{1}{p} g(t)v(t) + k(t, t)v(t) + \int_{\alpha}^t k_t(t, \sigma) v(\sigma) d\sigma \end{aligned}$$

then

$$v'(t) \leq A_6(t) + B_6(t)v(t) \quad \dots\dots\dots (2.44)$$

**Retarded Integral Inequalities with Iterated Integrals**

Ali W.K. Sangawi<sup>1,2</sup> and Sudad M. Rasheed<sup>1,2</sup>

Integrating both sides of (2.44) from  $\alpha$  to  $t$ , for  $t \in I$ , and using (2.42), we get:

$$z(t) \leq v(\alpha) \exp^{\int_{\alpha}^t R_2(\xi) d\xi} + \int_{\alpha}^t A_6(\eta) \exp^{\int_{\eta}^t B_2(\xi) d\xi} d\eta$$

..... (2.45)

from (2.45) and (2.41), one can get:

$$v(\alpha) \left[ 1 - \int_{\alpha}^{\beta} r(\sigma) \exp^{\int_{\alpha}^{\sigma} B_2(\xi) d\xi} d\sigma \right] \leq \int_{\alpha}^{\beta} r(\sigma) \int_{\alpha}^{\sigma} A_6(\eta) \exp^{\int_{\eta}^{\sigma} B_2(\xi) d\xi} d\eta d\sigma$$

from (2.20) which implies

$$v(\alpha) \leq M_3 \quad \text{..... (2.46)}$$

The required inequality (2.21) follows from (2.46), (2.45) and (2.9). From the hypotheses, we observe that  $\alpha'(t) \geq 0$  for  $t \in I_1$ .

**Iterated Retarded Integral Inequalities**

In this section we prove obtain explicit bounds to unknown functions in the some iterated retarded integral inequalities ,in the following theorem we take the single integral inequalities and in another theorem we take the double and triple integral inequalities.

**Theorem : (3.1)**

Let  $u(t), f(t), a(t), g(t)$  and  $h(t) \in C(I, \mathbb{R}_+)$ ,  $k(t, s) \in C(I^2, \mathbb{R}_+)$  for  $t, s \leq t \leq T$ ,  $\alpha(t) \in C^1(I, I)$  be non-decreasing with  $\alpha(t) \leq t$  on  $I$  and  $p > 1$  be real constant.

(c<sub>1</sub>)

If

Retarded Integral Inequalities with Iterated Integrals

Ali W.K. Sangawi<sup>1,2</sup> and Sudad M. Rasheed<sup>1,2</sup>

$$u^p(t) \leq a(t) + \int_{\alpha(t)}^t f(s) \left\{ u^p(s) + \int_{\alpha(s)}^s k(s, \nu) u(\nu) d\nu \right\} ds \quad \dots\dots (3.1)$$

for  $t \in I$ , then

$$u(t) \leq \left[ a(t) + \int_{\alpha(t)}^t D_1(s) \exp \int_s^{\alpha(t)} E_1(\xi) d\xi ds \right]^{\frac{1}{p}}, \quad \dots\dots (3.2)$$

for  $t \in I$ , where

$$D_1(t) = f(t) \left[ a(t) + \int_{\alpha(t)}^t k(t, \sigma) \left[ \frac{p-1}{p} + \frac{\alpha(\sigma)}{p} \right] d\sigma \right] \quad \text{and}$$

$$E_1(t) = f(t) \left[ 1 + \int_{\alpha(t)}^t \frac{k(t, \sigma)}{p} d\sigma \right].$$

(c<sub>2</sub>)

Let  $c(t)$  be real-valued positive continuous and nondecreasing function defined in  $I$ .

If

$$c^p(t) < c^p(t) + \int_{\alpha(t)}^t f(s) \left\{ u^p(s) + \int_{\alpha(s)}^s k(s, \sigma) u(\sigma) d\sigma \right\} ds, \quad \dots\dots (3.3)$$

for  $t \in I$ , then

$$u(t) \leq c(t) \left[ 1 + \int_{\alpha(t)}^t D_2(s) \exp \int_s^{\alpha(t)} E_2(\xi) d\xi ds \right]^{\frac{1}{p}} \quad \dots\dots (3.4)$$

for  $t \in I$ , where

$$D_2(t) = f(t) \left[ 1 + \int_{\alpha(t)}^t k(t, \sigma) c^{1-p}(\sigma) d\sigma \right] \quad \text{and}$$

$$E_2(t) = f(t) \left[ 1 + \int_{\alpha(t)}^t \frac{k(t, \sigma) c^{1-p}(\sigma)}{p} d\sigma \right].$$

**Proof :**

(c<sub>1</sub>)

Retarded Integral Inequalities with Iterated Integrals

Ali W.K. Sangawi<sup>1,2</sup> and Sudad M. Rasheed<sup>1,2</sup>

Define a function  $z(t)$  by  $z(t) = \int_{\alpha(t_*)}^{\alpha(t)} f(s) \left\{ u^p(s) + \int_{\alpha(t_*)}^s k(s, \sigma) u(\sigma) d\sigma \right\} ds$ .

..... (3.5)

Then  $z(t_*) = 0$  and as in the proof of part (a<sub>1</sub>), we get

$$z'(t) \leq \left[ f(\alpha(t)) \left\{ \alpha(\alpha(t)) + \int_{\alpha(t_*)}^{\alpha(t)} k(\alpha(t), \sigma) \left[ \frac{p-1}{p} + \frac{u(\sigma)}{p} \right] d\sigma \right\} \alpha'(t) + f(\alpha(t)) \left\{ 1 + \int_{\alpha(t_*)}^{\alpha(t)} \frac{k(\alpha(t), \sigma)}{p} d\sigma \right\} \alpha'(t) z(t) \right]$$

Therefore,  $t_* \leq \eta \leq t \leq T$ , one can have:

$$\frac{d}{d\eta} \left[ z(\eta) \exp^{\int_{\eta}^t f(\alpha(\tau)) \left\{ 1 + \int_{\alpha(t_*)}^{\alpha(\tau)} \frac{k(\alpha(\tau), \sigma)}{p} d\sigma \right\} \alpha'(\tau) d\tau} \right] \leq f(\alpha(\eta)) \left\{ \alpha(\alpha(\eta)) \int_{\alpha(t_*)}^{\alpha(\eta)} k(\alpha(\eta), \sigma) \left[ \frac{p-1}{p} + \frac{u(\sigma)}{p} \right] d\sigma + \alpha'(\eta) \exp^{\int_{\eta}^t f(\alpha(\tau)) \left\{ 1 + \int_{\alpha(t_*)}^{\alpha(\tau)} \frac{k(\alpha(\tau), \sigma)}{p} d\sigma \right\} \alpha'(\tau) d\tau} \right]$$

Integrating both side of the above inequality from  $t_*$  to  $t$ ,  $t \in I$  and by making the change of variables we get:

$$z(t) \leq \int_{\alpha(t_*)}^{\alpha(t)} D_1(s) \exp^{\int_s^{\alpha(t)} E_1(\xi) d\xi} ds \quad \text{for } t \in I \dots (3.6)$$

Using (3.6) in (2.8), yields the required inequality in (3.2).u

(c<sub>2</sub>)

Retarded Integral Inequalities with Iterated Integrals

Ali W.K. Sangawi<sup>1,2</sup> and Sudad M. Rasheed<sup>1,2</sup>

Since  $c(t)$  is positive continuous and nondecreasing function for  $t \in I$ , then inequality (3.3) can be written as

$$\left[ \frac{u(t)}{c(t)} \right]^p \leq 1 + \int_{\alpha(t_0)}^{\alpha(t)} f(s) \left\{ \left[ \frac{u(s)}{c(s)} \right]^p + \int_{\alpha(t_0)}^s k(s, \sigma) c^{1-p}(\sigma) \left[ \frac{u(\sigma)}{c(\sigma)} \right] d\sigma \right\} ds .$$

Now an application of the inequality given in  $(c_1)$  yields desired result in (3.4).

**Theorem : (3.2)**

Let

$u(t), g(t) \in C(I_1, \mathbb{R}_+)$ ,  $k(t, s), b(t, s), c(t, s) \in C(D_1, \mathbb{R}_+)$ ,  $h(t, s, \sigma) \in C(E_2, \mathbb{R}_+)$  and  $a(t), a'(t) \in C(I_1, \mathbb{R}_+)$ ,  $p > 1$  be a real constant,  $\alpha(t) \in C^1(I_1, I_1)$  be nondecreasing with  $\alpha(t) < t$  on  $I_1$ .

$(d_1)$

Let  $\phi(t) \in C(I_1, \mathbb{R}_+)$  and

$$\begin{aligned} u^p(t) \leq & a(t) + \int_{\alpha(t_0)}^{\alpha(t)} \phi(s)u(s)ds \\ & + \int_{\alpha(t_0)}^{\alpha(t)} \int_{\alpha(t_0)}^s k(s, \tau)u(\tau)d\tau ds \\ & + \int_{\alpha(t_0)}^{\alpha(t)} \int_{\alpha(t_0)}^s \int_{\alpha(t_0)}^{\tau} h(s, \tau, \sigma)u(\sigma)d\sigma d\tau ds \\ & + \int_{\alpha(t_0)}^{\beta} \int_{\alpha(t_0)}^s c(s, \tau)u(\tau)d\tau ds, \end{aligned} \dots (3.7)$$

for  $t \in I_1$ . If

$$q_1 = \frac{1}{p} \int_{\alpha(t_0)}^{\beta} \int_{\alpha(t_0)}^s c(s, \tau) \exp^{\int_{\alpha(t_0)}^{\alpha(\tau)} E_2(\theta) a \theta} d\tau ds < 1 \dots (3.8)$$

then

Retarded Integral Inequalities with Iterated Integrals

Ali W.K. Sangawi<sup>1,2</sup> and Sudad M. Rasheed<sup>1,2</sup>

$$u(t) \leq \left[ a(t) + N_1 \exp^{\int_{\alpha(t)}^t E_3(\theta) d\theta} + \int_{\alpha(t)}^t D_3(\psi) \exp^{\int_{\psi}^{\alpha(\tau)} E_3(\theta) d\theta} d\psi \right]^p,$$

for  $t \in I_1$ , where

$$D_3(t) = \frac{1}{p} \left\{ \phi(t)[(p-1) + a(t)] + \int_{\alpha(t)}^t k(t, \tau) [(p-1) + a(\tau)] d\tau + \int_{\alpha(t)}^t \int_{\alpha(\tau)}^{\tau} h(t, \tau, \sigma) [(p-1) + a(\sigma)] d\sigma d\tau \right\},$$

$$E_3(t) = \frac{1}{p} \left\{ \phi(t) + \int_{\alpha(t)}^t k(t, \tau) d\tau + \int_{\alpha(t)}^t \int_{\alpha(\tau)}^{\tau} h(t, \tau, \sigma) d\sigma d\tau \right\},$$

and

$$N_1 = \frac{1}{1 - q_1} \left\{ \frac{1}{p} \int_{\alpha(t)}^{\beta} \int_{\alpha(\tau)}^{\tau} c(s, \tau) \left[ (p-1) + a(\tau) + \int_{\alpha(\tau)}^{\tau} D_3(\psi) \exp^{\int_{\psi}^{\alpha(\tau)} E_3(\theta) d\theta} d\psi \right] d\tau ds \right\}$$

(d<sub>2</sub>)

Let  $k(t, s)$ ,  $b(t, s)$ ,  $h(t, s, \sigma)$  are nondecreasing in  $t \in I_1$ , for each  $s \in I_1$  and

Retarded Integral Inequalities with Iterated Integrals

Ali W.K. Sangawi<sup>1,2</sup> and Sudad M. Rasheed<sup>1,2</sup>

$$\begin{aligned}
 u^p(t) &\leq a(t) + \int_{\alpha(t^-)}^{\alpha(t)} k(t, \tau) u(\tau) d\tau \\
 &+ \int_{\alpha(t^-)}^{\alpha(t)} \int_{\alpha(\tau^-)}^{\tau} h(t, s, \sigma) u(\sigma) d\sigma ds + \\
 &\int_{\alpha(t^-)}^{\beta} b(t, s) \int_{\alpha(\tau^-)}^{\tau} c(s, \tau) u(\tau) d\tau ds,
 \end{aligned}
 \tag{3.10}$$

For  $t \in I_1$ . If

$$q_2 = \frac{1}{p} \int_{\alpha(t^-)}^{\beta} b(t, s) \int_{\alpha(\tau^-)}^{\tau} c(s, \tau) \exp^{\int_{\alpha(\tau^-)}^{\alpha(\tau)} E_4(\theta) d\theta} d\tau ds < 1,$$

..... (3.11)

then

$$u(t) \leq \left[ a(t) + N_2 \exp^{\int_{\alpha(t^-)}^{\alpha(t)} E_4(\theta) d\theta} + \int_{\alpha(t^-)}^{\alpha(t)} D_4(\psi) \exp^{\int_{\psi}^{\alpha(t)} E_4(\theta) d\theta} d\psi \right]^{\frac{1}{p}}$$

..... (3.12)

for  $t \in I_1$ , where

$$D_4(\psi) = \frac{1}{p} \left\{ k(t, \psi) [(p-1) + a(\psi)] + \int_{\alpha(\tau^-)}^{\psi} h(t, \psi, \sigma) [(p-1) + a(\sigma)] d\sigma \right\},$$

$$E_4(\theta) = \frac{1}{p} \left\{ k(t, \theta) + \int_{\alpha(\tau^-)}^{\theta} h(t, \theta, \sigma) d\sigma \right\}$$

and

Retarded Integral Inequalities with Iterated Integrals

Ali W.K. Sangawi<sup>1,2</sup> and Sudad M. Rasheed<sup>1,2</sup>

$$N_2 = \frac{1}{1 - q_2} \left[ \frac{1}{p} \int_{\alpha(t_2)}^{\beta} b(t, s) \int_{\alpha(t_2)}^s c(s, \tau) \left[ (p - 1) + a(\tau) + \int_{\alpha(\tau)}^{\tau} D_4(\psi) \exp\left\{ \int_{\psi}^{\alpha(\tau)} E_4(\theta) d\theta \right\} d\psi \right] d\tau ds \right].$$

(d<sub>3</sub>)

Let  $e(t, s) \in C(D_1, \mathbb{R}_+)$  and if

$$u^p(t) \leq a(t) + \int_{\alpha(t_2)}^{\alpha(t)} g(s) \left\{ u(s) + \int_{\alpha(t_2)}^s k(s, \sigma) u(\sigma) d\sigma + \int_{\alpha(t_2)}^{\beta} e(s, \sigma) u(\sigma) d\sigma \right\} ds, \tag{3.13}$$

for  $t \in I_1$ . Then

$$u(t) \leq \left[ a(t) + \int_{\alpha(t_2)}^{\alpha(t)} D_5(\psi) \exp\left\{ \int_{\psi}^{\alpha(t)} E_5(\theta) d\theta \right\} d\psi \right]^{\frac{1}{p}} \tag{3.14}$$

For  $t \in I_1$ , where

$$D_5(t) = \frac{1}{p} g(t) \left\{ (p - 1) + a(t) + \int_{\alpha(t_2)}^t k(t, \sigma) [(p - 1) + a(\sigma)] d\sigma + \int_{\alpha(t_2)}^{\beta} e(t, \sigma) [(p - 1) + a(\sigma)] d\sigma \right\},$$

and

$$E_5(t) = \frac{1}{p} g(t) \left[ 1 + k(t, \sigma) d\sigma + \int_{\alpha(t_2)}^{\beta} e(t, \sigma) d\sigma \right].$$

**Proof:**

(d<sub>1</sub>)

Define a function  $z(t)$  by



Retarded Integral Inequalities with Iterated Integrals

Ali W.K. Sangawi<sup>1,2</sup> and Sudad M. Rasheed<sup>1,2</sup>

$$\begin{aligned}
 z(t) = & \int_{\alpha(t-)}^{\alpha(t)} \phi(s)u(s)ds + \int_{\alpha(t-)}^{\alpha(t)} \int_{\alpha(t-)}^s k(s, \tau)u(\tau)d\tau ds \\
 & + \int_{\alpha(t-)}^{\alpha(t)} \int_{\alpha(t-)}^s \int_{\alpha(t-)}^{\tau} h(s, \tau, \sigma)u(\sigma)d\sigma d\tau ds \\
 & + \int_{\alpha(t-)}^{\beta} \int_{\alpha(t-)}^s c(s, \tau)u(\tau)d\tau ds, \quad \dots\dots\dots (3.15)
 \end{aligned}$$

Then  $z(t) \geq 0$ ,  $z(t)$  is nondecreasing for  $t \in I_1$

$$z(t) = \int_{\alpha(t-)}^{\beta} \int_{\alpha(t-)}^s c(s, \tau)u(\tau)d\tau ds \quad \dots\dots\dots (3.16)$$

Then as in the proof of part (a<sub>1</sub>), from (3.15) we see that the inequalities (2.8) and (2.9) hold.

Differentiating (3.15) and using (2.9) and the fact that  $z(t)$  is nondecreasing in  $t$ , we get:

$$\begin{aligned}
 z'(t) = & \phi(\alpha(t))u(\alpha(t))\alpha'(t) + \int_{\alpha(t-)}^{\alpha(t)} k(\alpha(t), \tau)u(\tau)d\tau \alpha'(t) \\
 & + \int_{\alpha(t-)}^{\alpha(t)} \int_{\alpha(t-)}^{\tau} h(\alpha(t), \tau, \sigma)u(\sigma)d\sigma d\tau \alpha'(t)
 \end{aligned}$$

Retarded Integral Inequalities with Iterated Integrals

Ali W.K. Sangawi<sup>1,2</sup> and Sudad M. Rasheed<sup>1,2</sup>

$$\begin{aligned} &\leq \frac{1}{p} \left\{ \phi(\alpha(t))[(p-1) + a(\alpha(t))] \right. \\ &\quad + \int_{\alpha(t^-)}^{\alpha(t)} k(\alpha(t), \tau) [(p-1) + a(\tau)] d\tau \\ &\quad + \int_{\alpha(t^-)}^{\alpha(t)} \int_{\alpha(t^-)}^{\tau} h(\alpha(t), \tau, \sigma) [(p-1) \\ &\quad \left. + a(\sigma)] d\sigma d\tau \right\} \alpha'(t) \\ &\quad + \frac{1}{p} \left\{ \phi(\alpha(t)) + \int_{\alpha(t^-)}^{\alpha(t)} k(\alpha(t), \tau) d\tau \right. \\ &\quad \left. + \int_{\alpha(t^-)}^{\alpha(t)} \int_{\alpha(t^-)}^{\tau} h(\alpha(t), \tau, \sigma) d\sigma d\tau \right\} \alpha'(t) z(t). \end{aligned}$$

Therefore,  $t_0 \leq \eta \leq t \leq \beta$ , one can have:

$$\begin{aligned} &\frac{d}{d\eta} \left\{ z(\eta) \exp^{\int_{\eta}^t \frac{1}{p} [\phi(\alpha(\xi)) + \int_{\alpha(\tau^-)}^{\alpha(\xi)} k(\alpha(\xi), \tau) d\tau + \int_{\alpha(\tau^-)}^{\alpha(\xi)} \int_{\alpha(\tau^-)}^{\tau} h(\alpha(\xi), \tau, \sigma) d\sigma d\tau] \alpha'(\xi) d\xi} \right\} \\ &\leq \frac{1}{p} \{ \phi(\alpha(\eta))[(p-1) + a(\alpha(\eta))] \\ &\quad + \int_{\alpha(\eta^-)}^{\alpha(\eta)} k(\alpha(\eta), \tau) [(p-1) + a(\tau)] d\tau \\ &\quad + \int_{\alpha(\eta^-)}^{\alpha(\eta)} \int_{\alpha(\eta^-)}^{\tau} h(\alpha(\eta), \tau, \sigma) [(p-1) \\ &\quad + a(\sigma)] d\sigma d\tau \} \\ &\quad \exp^{\int_{\eta}^t \frac{1}{p} [\phi(\alpha(\xi)) + \int_{\alpha(\tau^-)}^{\alpha(\xi)} k(\alpha(\xi), \tau) d\tau + \int_{\alpha(\tau^-)}^{\alpha(\xi)} \int_{\alpha(\tau^-)}^{\tau} h(\alpha(\xi), \tau, \sigma) d\sigma d\tau] \alpha'(\xi) d\xi} \alpha'(\eta) \end{aligned}$$

Retarded Integral Inequalities with Iterated Integrals

Ali W.K. Sangawi<sup>1,2</sup> and Sudad M. Rasheed<sup>1,2</sup>

Integrating both sides of the above inequality from  $t_*$  to  $t$ ,  $t \in I_1$  and by making the change of variables, we get:

$$z(t) \leq z(t_*) \exp^{\int_{\alpha(t_*)}^{\alpha(t)} E_z(\theta) d\theta} + \int_{\alpha(t_*)}^{\alpha(t)} D_3(\psi) \exp^{\int_{\psi}^{\alpha(t)} E_z(\theta) d\theta} d\psi \quad \dots (3.17)$$

from (3.17), (2.9) and (3.16), we get:

$$z(t) \leq \int_{\alpha(t_*)}^{\beta} \int_{\alpha(\tau)}^s c(s, \tau) \frac{1}{p} [(p-1) + \alpha(\tau)] + z(t_*) \exp^{\int_{\alpha(t_*)}^{\alpha(\tau)} E_z(\theta) d\theta} + \int_{\alpha(t_*)}^{\alpha(\tau)} D_3(\psi) \exp^{\int_{\psi}^{\alpha(\tau)} E_z(\theta) d\theta} d\psi \Big] d\tau ds,$$

then

$$z(t) \left[ 1 - \frac{1}{p} \int_{\alpha(t_*)}^{\beta} \int_{\alpha(\tau)}^s c(s, \tau) \exp^{\int_{\alpha(\tau)}^{\alpha(t)} E_z(\theta) d\theta} d\tau ds \right] \leq \int_{\alpha(t_*)}^{\beta} \int_{\alpha(\tau)}^s c(s, \tau) \frac{1}{p} \left[ (p-1) + \alpha(\tau) + \int_{\alpha(t_*)}^{\alpha(\tau)} D_3(\psi) \exp^{\int_{\psi}^{\alpha(\tau)} E_z(\theta) d\theta} d\psi \right] d\tau ds .$$

from (3.8) we obtain that

$$z(t_*) \leq N_1 . \quad \dots\dots\dots (3.18)$$

The required inequality (3.9) follows from (3.18), (3.17) and (2.8).

(d<sub>2</sub>)

Fix any  $T$ ,  $t_* \leq T \leq \beta$ , then for  $t_* \leq t \leq T$ , we have

Retarded Integral Inequalities with Iterated Integrals

Ali W.K. Sangawi<sup>1,2</sup> and Sudad M. Rasheed<sup>1,2</sup>

$$\begin{aligned}
 u^p(t) \leq & a(t) + \int_{\alpha(t_0)}^{\alpha(t)} k(T, \tau) u(\tau) d\tau \\
 & + \int_{\alpha(t_0)}^{\alpha(t)} \int_{\alpha(t_0)}^s h(T, s, \sigma) u(\sigma) d\sigma ds \\
 & + \int_{\alpha(t_0)}^{\beta} b(T, s) \int_{\alpha(t_0)}^s c(s, \tau) u(\tau) d\tau ds
 \end{aligned}$$

.....(3.19)

Define a function  $z(t, T)$ ,  $t_0 \leq t \leq T$  by

$$\begin{aligned}
 z(t, T) = & \int_{\alpha(t_0)}^{\alpha(t)} k(T, \tau) u(\tau) d\tau + \int_{\alpha(t_0)}^{\alpha(t)} \int_{\alpha(t_0)}^s h(T, s, \sigma) u(\sigma) d\sigma ds + \\
 & \int_{\alpha(t_0)}^{\beta} b(T, s) \int_{\alpha(t_0)}^s c(s, \tau) u(\tau) d\tau ds
 \end{aligned}$$

.....(3.20)

Then  $z(t, T) \leq 0$ ,  $z(t, T)$  is nondecreasing for  $t \in I_1$ ,

$$z(t_0, T) = \int_{\alpha(t_0)}^{\beta} b(T, s) \int_{\alpha(t_0)}^s c(s, \tau) u(\tau) d\tau ds, \quad \dots (3.21)$$

and inequality (3.10) can be written as

$$u^p(t) \leq a(t) + z(t, T), \quad t_0 \leq t \leq T \quad \dots(3.22)$$

Then as in the proof of part (a<sub>1</sub>), from (3.22) we see that the inequalities (3.23) hold.

$$u(t) \leq \frac{(p-1)}{p} + \frac{a(t)}{p} + \frac{z(t, T)}{p}, \quad t_0 \leq t \leq T \quad \dots(3.23)$$

Differentiating (3.20) and using (3.23) and the fact that  $z(t, T)$  is nondecreasing in  $t$ , we get:

$$\begin{aligned}
 z'(t, T) = & k(T, \alpha(t)) u(\alpha(t)) \alpha'(t) \\
 & + \int_{\alpha}^t h(T, \alpha(t), \sigma) u(\sigma) d\sigma \alpha'(t)
 \end{aligned}$$

Retarded Integral Inequalities with Iterated Integrals

Ali W.K. Sangawi<sup>1,2</sup> and Sudad M. Rasheed<sup>1,2</sup>

$$\begin{aligned} &\leq \frac{1}{p} \left\{ k(T, \alpha(t)) [(p-1) + a(\alpha(t))] \right. \\ &\quad \left. + \int_{\alpha(t_*)}^{\alpha(t)} h(T, \alpha(t), \sigma) [(p-1) + a(\sigma)] d\sigma \right\} \alpha'(t) \\ &\quad + \frac{1}{p} \left\{ k(T, \alpha(t)) + \int_{\alpha(t_*)}^{\alpha(t)} h(T, \alpha(t), \sigma) d\sigma \right\} \alpha'(t) z(t, T), \end{aligned}$$

..... (3.24)

for  $t_* < T$  by setting  $t = \xi$  in (3.24) and integrating it with respect to  $\xi$  from  $t_*$  to  $T$  and by making change of variables we get:

$$\begin{aligned} &z(T, T) \\ &\leq z(t_*, T) \exp^{\int_{\alpha(t_*)}^{\alpha(T)} \frac{1}{p} \left\{ k(T, \theta) + \int_{\alpha(t_*)}^{\theta} h(T, \theta, \sigma) d\sigma \right\} d\theta} \\ &\quad + \int_{\alpha(t_*)}^{\alpha(T)} \frac{1}{p} \left\{ k(T, \psi) [(p-1) + a(\psi)] \right. \\ &\quad \left. + \int_{\alpha(t_*)}^{\psi} h(T, \psi, \sigma) [(p-1) + a(\sigma)] d\sigma \right\} \\ &\quad \exp^{\int_{\alpha(t_*)}^{\alpha(T)} \frac{1}{p} \left\{ k(T, \theta) + \int_{\alpha(t_*)}^{\theta} h(T, \theta, \sigma) d\sigma \right\} d\theta} d\psi. \end{aligned}$$

Since  $T$  is arbitrary from (3.25), (3.23), (3.22) and (3.21) with  $T$  replaced by  $t$ , one can get:

$$\begin{aligned} &z(t, t) \leq z(t_*, t) \exp^{\int_{\alpha(t_*)}^{\alpha(t)} \frac{1}{p} \left\{ k(t, \theta) + \int_{\alpha(t_*)}^{\theta} h(t, \theta, \sigma) d\sigma \right\} d\theta} + \\ &\int_{\alpha(t_*)}^{\alpha(t)} \frac{1}{p} \left\{ k(t, \psi) [(p-1) + a(\psi)] + \int_{\alpha(t_*)}^{\psi} h(t, \psi, \sigma) [(p-1) + \right. \\ &\left. a(\sigma)] d\sigma \right\} \exp^{\int_{\alpha(t_*)}^{\alpha(t)} \frac{1}{p} \left\{ k(t, \theta) + \int_{\alpha(t_*)}^{\theta} h(t, \theta, \sigma) d\sigma \right\} d\theta} d\psi, \end{aligned}$$

.....(3.26)

Retarded Integral Inequalities with Iterated Integrals

Ali W.K. Sangawi<sup>1,2</sup> and Sudad M. Rasheed<sup>1,2</sup>

$$u^p(t) \leq a(t) + z(t, t) , \quad \dots\dots\dots (3.27)$$

and

$$u(t) \leq \frac{1}{p} \{ (p-1) + a(t) + z(t, t) \} \leq$$

$$\frac{1}{p} \left\{ (p-1) + a(t) + z(t, t) \exp^{\int_{\alpha(t)}^{\alpha(t)} E_4(\theta) d\theta} + \int_{\alpha(t)}^{\alpha(t)} D_4(\psi) \exp^{\int_{\psi}^{u(t)} E_4(\theta) d\theta} d\psi \right\}, \quad \dots\dots\dots(3.28)$$

and

$$z(t, t) = \int_{\alpha(t)}^{\beta} b(t, s) \int_{\alpha(s)}^s c(s, \tau) u(\tau) d\tau ds \quad \dots\dots\dots(3.29)$$

from (3.29) and (3.28) one can get:

$$\begin{aligned} z(t, t) &= \int_{\alpha(t)}^{\beta} b(t, s) \int_{\alpha(s)}^s c(s, \tau) \frac{1}{p} \left\{ (p-1) + a(\tau) \right. \\ &\quad \left. + z(t, \tau) \exp^{\int_{\alpha(\tau)}^{\alpha(\tau)} E_4(\theta) d\theta} \right. \\ &\quad \left. + \int_{\alpha(\tau)}^{\alpha(\tau)} D_4(\psi) \exp^{\int_{\psi}^{u(\tau)} E_4(\theta) d\theta} d\psi \right\} d\tau ds. \end{aligned}$$

Since  $z(t, T)$  is nondecreasing and nonnegative for  $t \in I_1$  and  $\tau \leq s \leq t \leq \beta$  then

Retarded Integral Inequalities with Iterated Integrals

Ali W.K. Sangawi<sup>1,2</sup> and Sudad M. Rasheed<sup>1,2</sup>

$$\begin{aligned}
 z(t, t) & \left[ 1 - \frac{1}{p} \int_{\alpha(t)}^{\beta} b(t, s) \int_{\alpha(t)}^s c(s, \tau) \exp^{\int_{\alpha(t)}^{\tau} E_4(\theta) d\theta} d\tau ds \right] \\
 & \leq \int_{\alpha(t)}^{\beta} b(t, s) \int_{\alpha(t)}^s c(s, \tau) \frac{1}{p} \{ (p-1) + a(\tau) \\
 & \quad + \int_{\alpha(t)}^{\tau} D_4(\psi) \exp^{\int_{\psi}^{\alpha(\tau)} E_4(\theta) d\theta} d\psi \} d\tau ds.
 \end{aligned}$$

From (3.11) which implies

$$z(t, t) \leq N_2 \dots\dots\dots(3.30)$$

The required inequality (3.12) follows from (3.30), (3.26), and (3.27).

(d<sub>3</sub>)

Define a function  $z(t)$  by

$$\begin{aligned}
 z(t) & = \int_{\alpha(t)}^{\alpha(t)} g(s) \left\{ u(s) + \int_{\alpha(t)}^s k(s, \sigma) u(\sigma) d\sigma + \right. \\
 & \quad \left. \int_{\alpha(t)}^{\beta} e(s, \sigma) u(\sigma) d\sigma \right\} ds \dots\dots\dots(3.31)
 \end{aligned}$$

Then  $z(t) \geq 0$ ,  $z(t)$  is nondecreasing for  $t \in I_1$ ,  $z(t_0) = 0$ .

Then as in the proof of part (a<sub>1</sub>), from (3.31) we see that the inequalities (2.8) and (2.9) hold.

Differentiating (3.31) and using (2.9) and the fact that  $z(t)$  is nondecreasing in  $t$ , we get:

$$z'(t) = g(\alpha(t))$$

Retarded Integral Inequalities with Iterated Integrals

Ali W.K. Sangawi<sup>1,2</sup> and Sudad M. Rasheed<sup>1,2</sup>

$$\left[ u(\alpha(t)) + \int_{\alpha(t_*)}^{\alpha(t)} k(\alpha(t), \sigma)u(\sigma) d\sigma + \int_{\alpha(t_*)}^{\beta} e(\alpha(t), \sigma)u(\sigma) d\sigma \right] \alpha'(t) \leq \frac{1}{p}g(\alpha(t))\{ (p-1) + a(\alpha(t)) + \int_{\alpha(t_*)}^{\alpha(t)} k(\alpha(t), \sigma) [(p-1) + a(\sigma)] d\sigma + \int_{\alpha(t_*)}^{\beta} e(\alpha(t), \sigma) [(p-1) + a(\sigma)] d\sigma \} \alpha'(t) + \frac{1}{p}g(\alpha(t)) \left\{ 1 + \int_{\alpha(t_*)}^{\alpha(t)} k(\alpha(t), \sigma) d\sigma + \int_{\alpha(t_*)}^{\beta} e(\alpha(t), \sigma) d\sigma \right\} \alpha'(t)z(t)$$

Therefore,  $t_* \leq \eta \leq t \leq \beta$ , one can have

$$\frac{d}{d\eta} \left\{ z(\eta) \exp^{\int_{\eta}^{t_1} \frac{1}{p}g(\alpha(\xi)) [1 + \int_{\alpha(t_*)}^{\alpha(\xi)} k(\alpha(\xi), \sigma) d\sigma + \int_{\alpha(t_*)}^{\beta} e(\alpha(\xi), \sigma) d\sigma] \alpha'(\xi) d\xi} \right\} \left\{ (p-1) + a(\alpha(\eta)) + \int_{\alpha(t_*)}^{\alpha(\eta)} k(\alpha(\eta), \sigma) [(p-1) + a(\sigma)] d\sigma + \int_{\alpha(t_*)}^{\beta} e(\alpha(\eta), \sigma) [(p-1) + a(\sigma)] d\sigma \right\} \exp^{\int_{\eta}^{t_1} \frac{1}{p}g(\alpha(\xi)) [1 + \int_{\alpha(t_*)}^{\alpha(\xi)} k(\alpha(\xi), \sigma) d\sigma + \int_{\alpha(t_*)}^{\beta} e(\alpha(\xi), \sigma) d\sigma] \alpha'(\xi) d\xi} \alpha'(\eta).$$

Integrating both sides of the above inequality from  $t_*$  to  $t$ ,  $t \in I_1$ , since  $z(t_*) = 0$ , and by making the change of variables we get:

$$z(t) \leq \int_{\alpha(t_*)}^{\alpha(t)} D_{\bar{\sigma}}(\psi) \exp^{\int_{\psi}^{\alpha(t)} E_{\bar{\sigma}}(\theta) d\theta} d\psi. \quad \dots (3.32)$$

The required inequality (3.14) follows from (3.32) and (2.9).



### Conclusions

We have constructed some iterated integral inequalities then extended to the iterated retarded integral inequalities. And also explicit bounds to unknown functions in each integral inequalities are given.

### References

1. Ali K., Several New Result on the Gronwall Delay Inequalities, *Kurdistan Academicians Journal*, 8(1)(2010)part A(107-114).
2. B.G. Pachpatte and Elsevier B.V., *Integral and Finite Difference Inequalities and Applications*, Amsterdam, Boston, London, New York, 2006.
3. B.G. Pachpatte, On a Certain Retarded Integral Inequality and its Applications, *J. Inequal. Pure and Appl. Math*, 5(1)(2004), Art.19.
4. B.G. Pachpatte, On Some New Nonlinear Retarded Integral Inequalities, *J. Inequal. Pure and Appl. math.*, 5(3)(2004), Art.80.
5. B.G. Pachpatte, Bounds on Certain Integral Inequalities, *J. Inequal. Pure and Appl. Math.*, 3(3)(2002), Art.47.
6. B.G. Pachpatte, Explicit Bounds on Certain Integral Inequalities, *J. Math. Anal. Appl.*, 267(2002), 48-61.
7. B.G. Pachpatte, *Inequalities for Finite Difference Equations*, Marcel, Dekker Inc., New York, 2002.
8. B.G. Pachpatte, On Some Retarded Integral Inequalities and Applications, *J. Inequal. Pure and Appl. Math.*, 3(2)(2002), Art.18.
9. B.G. Pachpatte, On Some New Inequalities Related to a Certain Inequality Arising in the Theory of Differential Equations, *J. Math. Anal. Appl.*, 251(2000), 736-751.
10. B.G. Pachpatte, *Inequalities for Differential and Integral Equation*, Academic Press, New York, 1998.
11. Dragomir S. S., *Some Gronwall type Inequalities and Application*, Academic Press, New York, 2003.

## Retarded Integral Inequalities with Iterated Integrals

Ali W.K. Sangawi<sup>1,2</sup> and Sudad M. Rasheed<sup>1,2</sup>

12. D.P. Mitrinovicetal and A. Fink, Classical and New Inequalities in Analysis, *Kluwer Academic Publishers, Dodrecht, Boston, London*, 1993.
13. D. Bainov and P. Simenove, Integral Inequalities and Applications, *Kluwer Academic Publish- ers, Dodrecht*, 1992.
14. D.S. Mitrinovic, Analytic Inequalities, *Springer-Verlag, Berlin, New York*, 1970.
15. Lakshmikantham V. and S. Leela, Differential and Integral Inequalities, *Vols. I, II, Academic Press, New York*, 1969.
16. Qing-Hua Ma and Josip Pecaric, On Some New Nonlinear Retarded Integral Inequalities with Iterated Integrals and Their Applications , *J. Korean Math.Soc.* 45(2008),2, 331-353.

